

Express Mail No. EL 998648560 US


**C10 ESTER SUBSTITUTED TAXANES****REFERENCE TO RELATED APPLICATION**

This application is a continuation of U.S. Patent Application No. 09/776,492 filed February 2, 2001 which claims priority from U.S. provisional application Serial No. 60/179,782, filed on February 2, 2000.

**BACKGROUND OF THE INVENTION**


The present invention is directed to novel taxanes which have exceptional utility as antitumor agents.

The taxane family of terpenes, of which baccatin III and taxol are members, has been the subject of considerable interest in both the biological and chemical arts. Taxol itself is employed as a cancer chemotherapeutic agent and possesses a broad range of tumor-inhibiting activity. Taxol has a 2'R, 3'S configuration and the following structural formula:



wherein Ac is acetyl.

Colin et al. reported in U.S. Patent 4,814,470 that certain taxol analogs have an activity significantly greater than that of taxol. One of these analogs, commonly referred to as docetaxel, has the following structural formula:



Express Mail No. EL 998648560 US

Although taxol and docetaxel are useful chemotherapeutic agents, there are limitations on their effectiveness, including limited efficacy against certain types of cancers and toxicity to subjects when administered at various doses. Accordingly, a need remains for additional chemotherapeutic agents with improved efficacy and less toxicity.

### SUMMARY OF THE INVENTION


Among the objects of the present invention, therefore, is the provision of taxanes which compare favorably to taxol and docetaxel with respect to efficacy as anti-tumor agents and with respect to toxicity. In general, these taxanes possess an ester substituent other than formate, acetate and heterosubstituted acetate at C-10, a hydroxy substituent at C-7 and a range of C-3' substituents.

Briefly, therefore, the present invention is directed to the taxane composition, per se, to pharmaceutical compositions comprising the taxane and a pharmaceutically acceptable carrier, and to methods of administration.

Other objects and features of this invention will be in part apparent and in part pointed out hereinafter.

### DETAILED DESCRIPTION OF THE PREFERRED EMBODIMENTS

In one embodiment of the present invention, the taxanes of the present invention correspond to structure (1):



(1)

wherein

R<sub>2</sub> is acyloxy;R<sub>7</sub> is hydroxy;R<sub>9</sub> is keto, hydroxy, or acyloxy;R<sub>10</sub> is R<sub>10a</sub>COO-;

Express Mail No. EL 998648560 US

$R_{10a}$  is hydrocarbyl, substituted hydrocarbyl, or heterocyclo wherein said hydrocarbyl or substituted hydrocarbyl contains carbon atoms in the alpha and beta positions relative to the carbon of which  $R_{10a}$  is a substituent;

$R_{14}$  is hydrido or hydroxy;

$X_3$  is substituted or unsubstituted alkyl, alkenyl, alkynyl, phenyl or heterocyclo, wherein alkyl comprises at least two carbon atoms;

$X_5$  is  $-\text{COX}_{10}$ ,  $-\text{COOX}_{10}$ , or  $-\text{CONHX}_{10}$ ;

$X_{10}$  is hydrocarbyl, substituted hydrocarbyl, or heterocyclo;

Ac is acetyl; and

$R_7$ ,  $R_9$ , and  $R_{10}$  independently have the alpha or beta stereochemical configuration.

In one embodiment,  $R_2$  is an ester ( $R_{2a}\text{C}(\text{O})\text{O}-$ ), a carbamate ( $R_{2a}R_{2b}\text{NC}(\text{O})\text{O}-$ ), a carbonate ( $R_{2a}\text{OC}(\text{O})\text{O}-$ ), or a thiocarbamate ( $R_{2a}\text{SC}(\text{O})\text{O}-$ ) wherein  $R_{2a}$  and  $R_{2b}$  are independently hydrogen, hydrocarbyl, substituted hydrocarbyl or heterocyclo. In a preferred embodiment,  $R_2$  is an ester ( $R_{2a}\text{C}(\text{O})\text{O}-$ ), wherein  $R_{2a}$  is aryl or heteroaromatic. In another preferred embodiment,  $R_2$  is an ester ( $R_{2a}\text{C}(\text{O})\text{O}-$ ), wherein  $R_{2a}$  is substituted or unsubstituted phenyl, furyl, thienyl, or pyridyl. In one particularly preferred embodiment,  $R_2$  is benzoyloxy.

While  $R_9$  is keto in one embodiment of the present invention, in other embodiments  $R_9$  may have the alpha or beta stereochemical configuration, preferably the beta stereochemical configuration, and may be, for example,  $\alpha$ - or  $\beta$ -hydroxy or  $\alpha$ - or  $\beta$ -acyloxy. For example, when  $R_9$  is acyloxy, it may be an ester ( $R_{9a}\text{C}(\text{O})\text{O}-$ ), a carbamate ( $R_{9a}R_{9b}\text{NC}(\text{O})\text{O}-$ ), a carbonate ( $R_{9a}\text{OC}(\text{O})\text{O}-$ ), or a thiocarbamate ( $R_{9a}\text{SC}(\text{O})\text{O}-$ ) wherein  $R_{9a}$  and  $R_{9b}$  are independently hydrogen, hydrocarbyl, substituted hydrocarbyl or heterocyclo. If  $R_9$  is an ester ( $R_{9a}\text{C}(\text{O})\text{O}-$ ),  $R_{9a}$  is substituted or unsubstituted alkyl, substituted or unsubstituted alkenyl, substituted or unsubstituted aryl or substituted or unsubstituted heteroaromatic. Still more preferably,  $R_9$  is an ester ( $R_{9a}\text{C}(\text{O})\text{O}-$ ), wherein  $R_{9a}$  is substituted or unsubstituted phenyl, substituted or unsubstituted furyl, substituted or unsubstituted thienyl, or substituted or unsubstituted pyridyl. In one embodiment  $R_9$  is ( $R_{9a}\text{C}(\text{O})\text{O}-$ ) wherein  $R_{9a}$  is methyl, ethyl, propyl (straight, branched or cyclic), butyl (straight, branched or cyclic), pentyl, (straight, branched or cyclic), or hexyl (straight, branched or cyclic). In another embodiment  $R_9$  is ( $R_{9a}\text{C}(\text{O})\text{O}-$ ) wherein  $R_{9a}$  is substituted methyl, substituted ethyl, substituted propyl (straight, branched or cyclic), substituted butyl (straight, branched or cyclic), substituted pentyl,

Express Mail No. EL 998648560 US

(straight, branched or cyclic), or substituted hexyl (straight, branched or cyclic) wherein the substituent(s) is/are selected from the group consisting of heterocyclo, alkoxy, alkenoxy, alkynoxy, aryloxy, hydroxy, protected hydroxy, keto, acyloxy, nitro, amino, amido, thiol, ketal, acetal, ester and ether moieties, but not phosphorous containing moieties.

In one embodiment,  $R_{10}$  is  $R_{10a}COO-$  wherein  $R_{10a}$  is (i) substituted or unsubstituted  $C_2$  to  $C_8$  alkyl (straight, branched or cyclic), such as ethyl, propyl, butyl, pentyl, or hexyl; (ii) substituted or unsubstituted  $C_2$  to  $C_8$  alkenyl (straight, branched or cyclic), such as ethenyl, propenyl, butenyl, pentenyl or hexenyl; (iii) substituted or unsubstituted  $C_2$  to  $C_8$  alkynyl (straight or branched) such as ethynyl, propynyl, butynyl, pentynyl, or hexynyl; (iv) substituted or unsubstituted phenyl; or (v) substituted or unsubstituted heteroaromatic such as furyl, thienyl, or pyridyl. The substituents may be hydrocarbonyl or any of the heteroatom containing substituents identified elsewhere herein for substituted hydrocarbonyl. In a preferred embodiment,  $R_{10a}$  is ethyl, straight, branched or cyclic propyl, straight, branched or cyclic butyl, straight, branched or cyclic pentyl, straight, branched or cyclic hexyl, straight or branched propenyl, isobutenyl, furyl or thienyl. In another embodiment,  $R_{10a}$  is substituted ethyl, substituted propyl (straight, branched or cyclic), substituted propenyl (straight or branched), substituted isobutenyl, substituted furyl or substituted thienyl wherein the substituent(s) is/are selected from the group consisting of heterocyclo, alkoxy, alkenoxy, alkynoxy, aryloxy, hydroxy, protected hydroxy, keto, acyloxy, nitro, amino, amido, thiol, ketal, acetal, ester and ether moieties, but not phosphorous containing moieties.


Exemplary  $X_3$  substituents include substituted or unsubstituted  $C_2$  to  $C_8$  alkyl, substituted or unsubstituted  $C_2$  to  $C_8$  alkenyl, substituted or unsubstituted  $C_2$  to  $C_8$  alkynyl, substituted or unsubstituted heteroaromatics containing 5 or 6 ring atoms, and substituted or unsubstituted phenyl. Exemplary preferred  $X_3$  substituents include substituted or unsubstituted ethyl, propyl, butyl, cyclopropyl, cyclobutyl, cyclohexyl, isobutenyl, furyl, thienyl, and pyridyl.

Exemplary  $X_5$  substituents include  $-COX_{10}$ ,  $-COOX_{10}$  or  $-CONHX_{10}$  wherein  $X_{10}$  is substituted or unsubstituted alkyl, alkenyl, phenyl or heteroaromatic. Exemplary preferred  $X_5$  substituents include  $-COX_{10}$ ,  $-COOX_{10}$  or  $-CONHX_{10}$  wherein  $X_{10}$  is (i) substituted or unsubstituted  $C_1$  to  $C_8$  alkyl such as substituted or unsubstituted methyl, ethyl, propyl (straight, branched or cyclic), butyl (straight, branched or cyclic), pentyl (straight, branched or cyclic), or hexyl (straight,

Express Mail No. EL 998648560 US

branched or cyclic); (ii) substituted or unsubstituted C<sub>2</sub> to C<sub>8</sub> alkenyl such as substituted or unsubstituted ethenyl, propenyl (straight, branched or cyclic), butenyl (straight, branched or cyclic), pentenyl (straight, branched or cyclic) or hexenyl (straight, branched or cyclic); (iii) substituted or unsubstituted C<sub>2</sub> to C<sub>8</sub> alkynyl such as substituted or unsubstituted ethynyl, propynyl (straight or branched), butynyl (straight or branched), pentynyl (straight or branched), or hexynyl (straight or branched); (iv) substituted or unsubstituted phenyl, or (v) substituted or unsubstituted heteroaromatic such as furyl, thienyl, or pyridyl, wherein the substituent(s) is/are selected from the group consisting of heterocyclo, alkoxy, alkenoxy, alkynoxy, aryloxy, hydroxy, protected hydroxy, keto, acyloxy, nitro, amino, amido, thiol, ketal, acetal, ester and ether moieties, but not phosphorous containing moieties.

In one embodiment of the present invention, the taxanes of the present invention correspond to structure (2):



(2)

wherein

R<sub>7</sub> is hydroxy;

R<sub>10</sub> is R<sub>10a</sub>COO-;

X<sub>3</sub> is substituted or unsubstituted alkyl, alkenyl, alkynyl, or heterocyclo,

wherein alkyl comprises at least two carbon atoms;

X<sub>5</sub> is -COX<sub>10</sub>, -COOX<sub>10</sub>, or -CONHX<sub>10</sub>; and

X<sub>10</sub> is hydrocarbyl, substituted hydrocarbyl, or heterocyclo; and

R<sub>10a</sub> is hydrocarbyl, substituted hydrocarbyl, or heterocyclo wherein said hydrocarbyl or substituted hydrocarbyl contains carbon atoms in the alpha and beta positions relative to the carbon of which R<sub>10a</sub> is a substituent;

Bz is benzoyl; and

Ac is acetyl.

Express Mail No. EL 998648560 US

For example, in this preferred embodiment in which the taxane corresponds to structure (2),  $R_{10a}$  may be substituted or unsubstituted ethyl, propyl or butyl, more preferably substituted or unsubstituted ethyl or propyl, still more preferably substituted or unsubstituted ethyl, and still more preferably unsubstituted ethyl.

5 While  $R_{10a}$  is selected from among these, in one embodiment  $X_3$  is selected from substituted or unsubstituted alkyl, alkenyl, phenyl or heterocyclo, more preferably substituted or unsubstituted alkenyl, phenyl or heterocyclo, still more preferably substituted or unsubstituted phenyl or heterocyclo, and still more preferably heterocyclo such as furyl, thienyl or pyridyl. While  $R_{10a}$  and  $X_3$  are selected from  
10 among these, in one embodiment  $X_5$  is selected from  $-COX_{10}$  wherein  $X_{10}$  is phenyl, alkyl or heterocyclo, more preferably phenyl. Alternatively, while  $R_{10a}$  and  $X_3$  are selected from among these, in one embodiment  $X_5$  is selected from  $-COX_{10}$  wherein  $X_{10}$  is phenyl, alkyl or heterocyclo, more preferably phenyl, or  $X_5$  is  $-COOX_{10}$  wherein  $X_{10}$  is alkyl, preferably t-butyl. Among the more preferred  
15 embodiments, therefore, are taxanes corresponding to structure 2 in which (i)  $X_5$  is  $-COOX_{10}$  wherein  $X_{10}$  is tert-butyl or  $X_5$  is  $-COX_{10}$  wherein  $X_{10}$  is phenyl, (ii)  $X_3$  is substituted or unsubstituted cycloalkyl, alkenyl, phenyl or heterocyclo, more preferably substituted or unsubstituted isobutenyl, phenyl, furyl, thienyl, or pyridyl, still more preferably unsubstituted isobutenyl, furyl, thienyl or pyridyl, and  
20 (iii)  $R_{7a}$  is unsubstituted ethyl or propyl, more preferably ethyl.

Among the preferred embodiments, therefore, are taxanes corresponding to structure 1 or 2 wherein  $R_{10}$  is  $R_{10a}COO^-$  wherein  $R_{10a}$  is ethyl. In this embodiment,  $X_3$  is preferably cycloalkyl, isobutenyl, or heterocyclo, more preferably heterocyclo, still more preferably furyl, thienyl or pyridyl; and  $X_5$  is  
25 preferably benzoyl, alkoxycarbonyl, or heterocyclocarbonyl, more preferably benzoyl, t-butoxycarbonyl or t-amylloxycarbonyl. In one alternative of this embodiment,  $X_3$  is heterocyclo;  $X_5$  is benzoyl, alkoxycarbonyl, or heterocyclocarbonyl, more preferably benzoyl, t-butoxycarbonyl or t-amylloxycarbonyl, still more preferably t-butoxycarbonyl;  $R_2$  is benzoyl,  $R_9$  is keto and  $R_{14}$  is hydrido. In another alternative of this embodiment,  $X_3$  is heterocyclo;  
30  $X_5$  is benzoyl, alkoxycarbonyl, or heterocyclocarbonyl, more preferably benzoyl, t-butoxycarbonyl or t-amylloxycarbonyl, still more preferably t-butoxycarbonyl;  $R_2$  is benzoyl,  $R_9$  is keto and  $R_{14}$  is hydrido. In another alternative of this embodiment,  $X_3$  is heterocyclo;  $X_5$  is benzoyl, alkoxycarbonyl, or heterocyclocarbonyl, more  
35 preferably benzoyl, t-butoxycarbonyl or t-amylloxycarbonyl, still more preferably t-

Express Mail No. EL 998648560 US


butoxycarbonyl;  $R_2$  is benzoyl,  $R_9$  is keto and  $R_{14}$  is hydroxy. In another alternative of this embodiment,  $X_3$  is heterocyclo;  $X_5$  is benzoyl, alkoxycarbonyl, or heterocyclocarbonyl, more preferably benzoyl, t-butoxycarbonyl or t-amylloxycarbonyl, still more preferably t-butoxycarbonyl;  $R_2$  is benzoyl,  $R_9$  is hydroxy and  $R_{14}$  is hydroxy. In another alternative of this embodiment,  $X_3$  is heterocyclo;  $X_5$  is benzoyl, alkoxycarbonyl, or heterocyclocarbonyl, more preferably benzoyl, t-butoxycarbonyl or t-amylloxycarbonyl, still more preferably t-butoxycarbonyl;  $R_2$  is benzoyl,  $R_9$  is hydroxy and  $R_{14}$  is hydrido. In another alternative of this embodiment,  $X_3$  is heterocyclo;  $X_5$  is benzoyl, alkoxycarbonyl, or heterocyclocarbonyl, more preferably benzoyl, t-butoxycarbonyl or t-amylloxycarbonyl, still more preferably t-butoxycarbonyl;  $R_2$  is benzoyl,  $R_9$  is acyloxy and  $R_{14}$  is hydroxy. In another alternative of this embodiment,  $X_3$  is heterocyclo;  $X_5$  is benzoyl, alkoxycarbonyl, or heterocyclocarbonyl, more preferably benzoyl, t-butoxycarbonyl or t-amylloxycarbonyl, still more preferably t-butoxycarbonyl;  $R_2$  is benzoyl,  $R_9$  is acyloxy and  $R_{14}$  is hydrido. In each of the alternatives of this embodiment when the taxane has structure 1,  $R_7$  and  $R_{10}$  may each have the beta stereochemical configuration,  $R_7$  and  $R_{10}$  may each have the alpha stereochemical configuration,  $R_7$  may have the alpha stereochemical configuration while  $R_{10}$  has the beta stereochemical configuration or  $R_7$  may have the beta stereochemical configuration while  $R_{10}$  has the alpha stereochemical configuration.

Also among the preferred embodiments are taxanes corresponding to structure 1 or 2 wherein  $R_{10}$  is  $R_{10a}COO^-$  wherein  $R_{10a}$  is propyl. In this embodiment,  $X_3$  is preferably cycloalkyl, isobutenyl, phenyl, substituted phenyl such as p-nitrophenyl, or heterocyclo, more preferably heterocyclo, still more preferably furyl, thienyl or pyridyl; and  $X_5$  is preferably benzoyl, alkoxycarbonyl, or heterocyclocarbonyl, more preferably benzoyl, t-butoxycarbonyl or t-amylloxycarbonyl. In one alternative of this embodiment,  $X_3$  is heterocyclo;  $X_5$  is benzoyl, alkoxycarbonyl, or heterocyclocarbonyl, more preferably benzoyl, t-butoxycarbonyl or t-amylloxycarbonyl, still more preferably t-butoxycarbonyl;  $R_2$  is benzoyl,  $R_9$  is keto and  $R_{14}$  is hydrido. In another alternative of this embodiment,  $X_3$  is heterocyclo;  $X_5$  is benzoyl, alkoxycarbonyl, or heterocyclocarbonyl, more preferably benzoyl, t-butoxycarbonyl or t-amylloxycarbonyl, still more preferably t-butoxycarbonyl;  $R_2$  is benzoyl,  $R_9$  is keto and  $R_{14}$  is hydrido. In another alternative of this embodiment,  $X_3$  is heterocyclo;  $X_5$  is benzoyl, alkoxycarbonyl, or

Express Mail No. EL 998648560 US

heterocyclocarbonyl, more preferably benzoyl, t-butoxycarbonyl or t-amyloxycarbonyl, still more preferably t-butoxycarbonyl;  $R_2$  is benzoyl,  $R_9$  is keto and  $R_{14}$  is hydroxy. In another alternative of this embodiment,  $X_3$  is heterocyclo;  $X_5$  is benzoyl, alkoxycarbonyl, or heterocyclocarbonyl, more preferably benzoyl, t-butoxycarbonyl or t-amyloxycarbonyl, still more preferably t-butoxycarbonyl;  $R_2$  is benzoyl,  $R_9$  is hydroxy and  $R_{14}$  is hydroxy. In another alternative of this embodiment,  $X_3$  is heterocyclo;  $X_5$  is benzoyl, alkoxycarbonyl, or heterocyclocarbonyl, more preferably benzoyl, t-butoxycarbonyl or t-amyloxycarbonyl, still more preferably t-butoxycarbonyl;  $R_2$  is benzoyl,  $R_9$  is hydroxy and  $R_{14}$  is hydrido. In another alternative of this embodiment,  $X_3$  is heterocyclo;  $X_5$  is benzoyl, alkoxycarbonyl, or heterocyclocarbonyl, more preferably benzoyl, t-butoxycarbonyl or t-amyloxycarbonyl, still more preferably t-butoxycarbonyl;  $R_2$  is benzoyl,  $R_9$  is acyloxy and  $R_{14}$  is hydroxy. In another alternative of this embodiment,  $X_3$  is heterocyclo;  $X_5$  is benzoyl, alkoxycarbonyl, or heterocyclocarbonyl, more preferably benzoyl, t-butoxycarbonyl or t-amyloxycarbonyl, still more preferably t-butoxycarbonyl;  $R_2$  is benzoyl,  $R_9$  is acyloxy and  $R_{14}$  is hydrido. In each of the alternatives of this embodiment when the taxane has structure 1,  $R_7$  and  $R_{10}$  may each have the beta stereochemical configuration,  $R_7$  and  $R_{10}$  may each have the alpha stereochemical configuration,  $R_7$  may have the alpha stereochemical configuration while  $R_{10}$  has the beta stereochemical configuration or  $R_7$  may have the beta stereochemical configuration while  $R_{10}$  has the alpha stereochemical configuration.

Taxanes having the general formula 1 may be obtained by treatment of a  $\beta$ -lactam with an alkoxide having the taxane tetracyclic nucleus and a C-13 metallic oxide substituent to form compounds having a  $\beta$ -amido ester substituent at C-13 (as described more fully in Holton U.S. Patent 5,466,834), followed by removal of the hydroxy protecting groups. The  $\beta$ -lactam has the following structural formula (3):




(3)



Express Mail No. EL 998648560 US

wherein  $P_2$  is a hydroxy protecting group and  $X_3$  and  $X_5$  are as previously defined and the alkoxide has the structural formula (4):



(4)

wherein M is a metal or ammonium,  $P_7$  is a hydroxy protecting group and  $R_{10}$  is as previously defined.

Alkoxide 4 may be prepared from 10-deacetylbaccatin III (or a derivative thereof) by selective protection of the C(7) hydroxyl group and then esterification of the C(10) hydroxyl group followed by treatment with a metallic amide. In one embodiment of the present invention, the C(7) hydroxyl group of 10-deacetylbaccatin III is selectively protected with a silyl group as described, for example, by Denis, et. al. (*J. Am. Chem. Soc.*, **1988**, 110, 5917). In general, the silylating agents may be used either alone or in combination with a catalytic amount of a base such as an alkali metal base.


Alternatively, the C(10) hydroxyl group of a taxane can be selectively acylated in the absence of a base, as described, for example in Holton et al., PCT Patent Application WO 99/09021. Acylating agents which may be used for the selective acylation of the C(10) hydroxyl group of a taxane include substituted or unsubstituted alkyl or aryl anhydrides. While the acylation of the C(10) hydroxy group of the taxane will proceed at an adequate rate for many acylating agents, it has been discovered that the reaction rate may be increased by including a Lewis acid in the reaction mixture. Preferred Lewis acids include zinc chloride, stannic chloride, cerium trichloride, cuprous chloride, lanthanum trichloride, dysprosium trichloride, and ytterbium trichloride. Zinc chloride or cerium trichloride is particularly preferred when the acylating agent is an anhydride.

Derivatives of 10-deacetylbaccatin III having alternative substituents at C(2), C(9) and C(14) and processes for their preparation are known in the art. Taxane derivatives having acyloxy substituents other than benzoyloxy at C(2) may be prepared, for example, as described in Holton et al., U.S. Patent No. 5,728,725 or Kingston et al., U.S. Patent No. 6,002,023. Taxanes having acyloxy

Express Mail No. EL 998648560 US

or hydroxy substituents at C(9) in place of keto may be prepared, for example as described in Holton et al., U.S. Patent No. 6,011,056 or Gunawardana et al., U.S. Patent No. 5,352,806. Taxanes having a beta hydroxy substituent at C(14) may be prepared from naturally occurring 14-hydroxy-10-deacetylbaccatin III.

Processes for the preparation and resolution of the  $\beta$ -lactam starting material are generally well known. For example, the  $\beta$ -lactam may be prepared as described in Holton, U.S. Patent No. 5,430,160 and the resulting enantiomeric mixtures of  $\beta$ -lactams may be resolved by a stereoselective hydrolysis using a lipase or enzyme as described, for example, in Patel, U.S. Patent No. 5,879,929 Patel U.S. Patent No. 5,567,614 or a liver homogenate as described, for example, in PCT Patent Application No. 00/41204. In a preferred embodiment in which the  $\beta$ -lactam is furyl substituted at the C(4) position, the  $\beta$ -lactam can be prepared as illustrated in the following reaction scheme:



Express Mail No. EL 998648560 US

wherein Ac is acetyl, NEt<sub>3</sub> is triethylamine, CAN is ceric ammonium nitrate, and p-TsOH is p-toluenesulfonic acid. The beef liver resolution may be carried out, for example, by combining the enantiomeric  $\beta$ -lactam mixture with a beef liver suspension (prepared, for example, by adding 20 g of frozen beef liver to a blender and then adding a pH 8 buffer to make a total volume of 1 L).

Compounds of formula 1 of the instant invention are useful for inhibiting tumor growth in mammals including humans and are preferably administered in the form of a pharmaceutical composition comprising an effective antitumor amount of a compound of the instant invention in combination with at least one pharmaceutically or pharmacologically acceptable carrier. The carrier, also known in the art as an excipient, vehicle, auxiliary, adjuvant, or diluent, is any substance which is pharmaceutically inert, confers a suitable consistency or form to the composition, and does not diminish the therapeutic efficacy of the antitumor compounds. The carrier is "pharmaceutically or pharmacologically acceptable" if it does not produce an adverse, allergic or other untoward reaction when administered to a mammal or human, as appropriate.

The pharmaceutical compositions containing the antitumor compounds of the present invention may be formulated in any conventional manner. Proper formulation is dependent upon the route of administration chosen. The compositions of the invention can be formulated for any route of administration so long as the target tissue is available via that route. Suitable routes of administration include, but are not limited to, oral, parenteral (e.g., intravenous, intraarterial, subcutaneous, rectal, subcutaneous, intramuscular, intraorbital, intracapsular, intraspinal, intraperitoneal, or intrasternal), topical (nasal, transdermal, intraocular), intravesical, intrathecal, enteral, pulmonary, intralymphatic, intracavitary, vaginal, transurethral, intradermal, aural, intramammary, buccal, orthotopic, intratracheal, intralesional, percutaneous, endoscopic, transmucosal, sublingual and intestinal administration.

Pharmaceutically acceptable carriers for use in the compositions of the present invention are well known to those of ordinary skill in the art and are selected based upon a number of factors: the particular antitumor compound used, and its concentration, stability and intended bioavailability; the disease, disorder or condition being treated with the composition; the subject, its age, size and general condition; and the route of administration. Suitable carriers are readily determined by one of ordinary skill in the art (see, for example, J. G. Nairn,

Express Mail No. EL 998648560 US

in: Remington's Pharmaceutical Science (A. Gennaro, ed.), Mack Publishing Co., Easton, Pa., (1985), pp. 1492-1517, the contents of which are incorporated herein by reference).

5 The compositions are preferably formulated as tablets, dispersible powders, pills, capsules, gelcaps, caplets, gels, liposomes, granules, solutions, suspensions, emulsions, syrups, elixirs, troches, dragees, lozenges, or any other dosage form which can be administered orally. Techniques and compositions for making oral dosage forms useful in the present invention are described in the following references: 7 Modern Pharmaceutics, Chapters 9 and 10 (Banker & Rhodes, Editors, 1979); Lieberman et al., Pharmaceutical Dosage Forms: Tablets (1981); and Ansel, Introduction to Pharmaceutical Dosage Forms 2nd Edition (1976).

15 The compositions of the invention for oral administration comprise an effective antitumor amount of a compound of the invention in a pharmaceutically acceptable carrier. Suitable carriers for solid dosage forms include sugars, starches, and other conventional substances including lactose, talc, sucrose, gelatin, carboxymethylcellulose, agar, mannitol, sorbitol, calcium phosphate, calcium carbonate, sodium carbonate, kaolin, alginic acid, acacia, corn starch, potato starch, sodium saccharin, magnesium carbonate, tragacanth, 20 microcrystalline cellulose, colloidal silicon dioxide, croscarmellose sodium, talc, magnesium stearate, and stearic acid. Further, such solid dosage forms may be uncoated or may be coated by known techniques; e.g., to delay disintegration and absorption.

25 The antitumor compounds of the present invention are also preferably formulated for parenteral administration, e.g., formulated for injection via intravenous, intraarterial, subcutaneous, rectal, subcutaneous, intramuscular, intraorbital, intracapsular, intraspinal, intraperitoneal, or intrasternal routes. The compositions of the invention for parenteral administration comprise an effective antitumor amount of the antitumor compound in a pharmaceutically acceptable carrier. Dosage forms suitable for parenteral administration include solutions, 30 suspensions, dispersions, emulsions or any other dosage form which can be administered parenterally. Techniques and compositions for making parenteral dosage forms are known in the art.

35 Suitable carriers used in formulating liquid dosage forms for oral or parenteral administration include nonaqueous, pharmaceutically-acceptable polar

Express Mail No. EL 998648560 US

solvents such as oils, alcohols, amides, esters, ethers, ketones, hydrocarbons and mixtures thereof, as well as water, saline solutions, dextrose solutions (e.g., DW5), electrolyte solutions, or any other aqueous, pharmaceutically acceptable liquid.

5           Suitable nonaqueous, pharmaceutically-acceptable polar solvents include, but are not limited to, alcohols (e.g.,  $\alpha$ -glycerol formal,  $\beta$ -glycerol formal, 1, 3-butylene glycol, aliphatic or aromatic alcohols having 2-30 carbon atoms such as methanol, ethanol, propanol, isopropanol, butanol, t-butanol, hexanol, octanol, amylene hydrate, benzyl alcohol, glycerin (glycerol), glycol, hexylene glycol, 10   tetrahydrofurfuryl alcohol, lauryl alcohol, cetyl alcohol, or stearyl alcohol, fatty acid esters of fatty alcohols such as polyalkylene glycols (e.g., polypropylene glycol, polyethylene glycol), sorbitan, sucrose and cholesterol); amides (e.g., dimethylacetamide (DMA), benzyl benzoate DMA, dimethylformamide, N-( $\beta$ -hydroxyethyl)-lactamide, N, N-dimethylacetamide, amides, 2-pyrrolidinone, 15   1-methyl-2-pyrrolidinone, or polyvinylpyrrolidone); esters (e.g., 1-methyl-2-pyrrolidinone, 2-pyrrolidinone, acetate esters such as monoacetin, diacetin, and triacetin, aliphatic or aromatic esters such as ethyl caprylate or octanoate, alkyl oleate, benzyl benzoate, benzyl acetate, dimethylsulfoxide (DMSO), esters of glycerin such as mono, di, or tri-glyceryl citrates or tartrates, ethyl benzoate, ethyl 20   acetate, ethyl carbonate, ethyl lactate, ethyl oleate, fatty acid esters of sorbitan, fatty acid derived PEG esters, glyceryl monostearate, glyceride esters such as mono, di, or tri-glycerides, fatty acid esters such as isopropyl myristate, fatty acid derived PEG esters such as PEG-hydroxyoleate and PEG-hydroxystearate, N-methyl pyrrolidinone, pluronic 60, polyoxyethylene sorbitol oleic polyesters such 25   as poly(ethoxylated)<sub>30-60</sub> sorbitol poly(oleate)<sub>2-4</sub>, poly(oxyethylene)<sub>15-20</sub> monooleate, poly(oxyethylene)<sub>15-20</sub> mono 12-hydroxystearate, and poly(oxyethylene)<sub>15-20</sub> mono ricinoleate, polyoxyethylene sorbitan esters such as polyoxyethylene-sorbitan monooleate, polyoxyethylene-sorbitan monopalmitate, polyoxyethylene-sorbitan monolaurate, polyoxyethylene-sorbitan monostearate, and Polysorbate® 20, 40, 30   60 or 80 from ICI Americas, Wilmington, DE, polyvinylpyrrolidone, alkyleneoxy modified fatty acid esters such as polyoxyl 40 hydrogenated castor oil and polyoxyethylated castor oils (e.g., Cremophor® EL solution or Cremophor® RH 40 solution), saccharide fatty acid esters (i.e., the condensation product of a monosaccharide (e.g., pentoses such as ribose, ribulose, arabinose, xylose, 35   lyxose and xylulose, hexoses such as glucose, fructose, galactose, mannose and

Express Mail No. EL 998648560 US

sorbose, trioses, tetroses, heptoses, and octoses), disaccharide (e.g., sucrose, maltose, lactose and trehalose) or oligosaccharide or mixture thereof with a C<sub>4</sub>-C<sub>22</sub> fatty acid(s)(e.g., saturated fatty acids such as caprylic acid, capric acid, lauric acid, myristic acid, palmitic acid and stearic acid, and unsaturated fatty acids such as palmitoleic acid, oleic acid, elaidic acid, erucic acid and linoleic acid)), or steroidal esters); alkyl, aryl, or cyclic ethers having 2-30 carbon atoms (e.g., diethyl ether, tetrahydrofuran, dimethyl isosorbide, diethylene glycol monoethyl ether); glycofurol (tetrahydrofurfuryl alcohol polyethylene glycol ether); ketones having 3-30 carbon atoms (e.g., acetone, methyl ethyl ketone, methyl isobutyl ketone); aliphatic, cycloaliphatic or aromatic hydrocarbons having 4-30 carbon atoms (e.g., benzene, cyclohexane, dichloromethane, dioxolanes, hexane, n-decane, n-dodecane, n-hexane, sulfolane, tetramethylenesulfon, tetramethylenesulfoxide, toluene, dimethylsulfoxide (DMSO), or tetramethylenesulfoxide); oils of mineral, vegetable, animal, essential or synthetic origin (e.g., mineral oils such as aliphatic or wax-based hydrocarbons, aromatic hydrocarbons, mixed aliphatic and aromatic based hydrocarbons, and refined paraffin oil, vegetable oils such as linseed, tung, safflower, soybean, castor, cottonseed, groundnut, rapeseed, coconut, palm, olive, corn, corn germ, sesame, persic and peanut oil and glycerides such as mono-, di- or triglycerides, animal oils such as fish, marine, sperm, cod-liver, haliver, squalene, squalane, and shark liver oil, oleic oils, and polyoxyethylated castor oil); alkyl or aryl halides having 1-30 carbon atoms and optionally more than one halogen substituent; methylene chloride; monoethanolamine; petroleum benzin; trolamine; omega-3 polyunsaturated fatty acids (e.g., alpha-linolenic acid, eicosapentaenoic acid, docosapentaenoic acid, or docosahexaenoic acid); polyglycol ester of 12-hydroxystearic acid and polyethylene glycol (Solutol® HS-15, from BASF, Ludwigshafen, Germany); polyoxyethylene glycerol; sodium laurate; sodium oleate; or sorbitan monooleate.

Other pharmaceutically acceptable solvents for use in the invention are well known to those of ordinary skill in the art, and are identified in The Chemotherapy Source Book (Williams & Wilkens Publishing), The Handbook of Pharmaceutical Excipients, (American Pharmaceutical Association, Washington, D.C., and The Pharmaceutical Society of Great Britain, London, England, 1968), Modern Pharmaceutics, (G. Banker et al., eds., 3d ed.)(Marcel Dekker, Inc., New York, New York, 1995), The Pharmacological Basis of Therapeutics, (Goodman &

Express Mail No. EL 998648560 US

Gilman, McGraw Hill Publishing), Pharmaceutical Dosage Forms, (H. Lieberman et al., eds., )(Marcel Dekker, Inc., New York, New York, 1980), Remington's Pharmaceutical Sciences (A. Gennaro, ed., 19th ed.)(Mack Publishing, Easton, PA, 1995), The United States Pharmacopeia 24, The National Formulary 19,  
5 (National Publishing, Philadelphia, PA, 2000), A.J. Spiegel et al., and Use of Nonaqueous Solvents in Parenteral Products, JOURNAL OF PHARMACEUTICAL SCIENCES, Vol. 52, No. 10, pp. 917-927 (1963).

Preferred solvents include those known to stabilize the antitumor compounds, such as oils rich in triglycerides, for example, safflower oil, soybean  
10 oil or mixtures thereof, and alkyleneoxy modified fatty acid esters such as polyoxyl 40 hydrogenated castor oil and polyoxyethylated castor oils (e.g., Cremophor® EL solution or Cremophor® RH 40 solution). Commercially available triglycerides include Intralipid® emulsified soybean oil (Kabi-Pharmacia Inc., Stockholm, Sweden), Nutralipid ® emulsion (McGaw, Irvine, California), Liposyn® II 20%  
15 emulsion (a 20% fat emulsion solution containing 100 mg safflower oil, 100 mg soybean oil, 12 mg egg phosphatides, and 25 mg glycerin per ml of solution; Abbott Laboratories, Chicago, Illinois), Liposyn® III 2% emulsion (a 2% fat emulsion solution containing 100 mg safflower oil, 100 mg soybean oil, 12 mg egg phosphatides, and 25 mg glycerin per ml of solution; Abbott Laboratories,  
20 Chicago, Illinois), natural or synthetic glycerol derivatives containing the docosahexaenoyl group at levels between 25% and 100% by weight based on the total fatty acid content (Dhasco® (from Martek Biosciences Corp., Columbia, MD), DHA Maguro® (from Daito Enterprises, Los Angeles, CA), Soyacal®, and Travemulsion®. Ethanol is a preferred solvent for use in dissolving the antitumor  
25 compound to form solutions, emulsions, and the like.

Additional minor components can be included in the compositions of the invention for a variety of purposes well known in the pharmaceutical industry. These components will for the most part impart properties which enhance  
30 retention of the antitumor compound at the site of administration, protect the stability of the composition, control the pH, facilitate processing of the antitumor compound into pharmaceutical formulations, and the like. Preferably, each of these components is individually present in less than about 15 weight % of the total composition, more preferably less than about 5 weight %, and most preferably less than about 0.5 weight % of the total composition. Some  
35 components, such as fillers or diluents, can constitute up to 90 wt.% of the total

Express Mail No. EL 998648560 US

composition, as is well known in the formulation art. Such additives include cryoprotective agents for preventing reprecipitation of the taxane, surface active, wetting or emulsifying agents (e.g., lecithin, polysorbate-80, Tween® 80, pluronic 60, polyoxyethylene stearate ), preservatives (e.g., ethyl-p-hydroxybenzoate), microbial preservatives (e.g., benzyl alcohol, phenol, m-cresol, chlorobutanol, sorbic acid, thimerosal and paraben), agents for adjusting pH or buffering agents (e.g., acids, bases, sodium acetate, sorbitan monolaurate), agents for adjusting osmolarity (e.g., glycerin), thickeners (e.g., aluminum monostearate, stearic acid, cetyl alcohol, stearyl alcohol, guar gum, methyl cellulose, hydroxypropylcellulose, tristearin, cetyl wax esters, polyethylene glycol), colorants, dyes, flow aids, non-volatile silicones (e.g., cyclomethicone), clays (e.g., bentonites), adhesives, bulking agents, flavorings, sweeteners, adsorbents, fillers (e.g., sugars such as lactose, sucrose, mannitol, or sorbitol, cellulose, or calcium phosphate), diluents (e.g., water, saline, electrolyte solutions), binders (e.g., starches such as maize starch, wheat starch, rice starch, or potato starch, gelatin, gum tragacanth, methyl cellulose, hydroxypropyl methylcellulose, sodium carboxymethyl cellulose, polyvinylpyrrolidone, sugars, polymers, acacia), disintegrating agents (e.g., starches such as maize starch, wheat starch, rice starch, potato starch, or carboxymethyl starch, cross-linked polyvinyl pyrrolidone, agar, alginic acid or a salt thereof such as sodium alginate, croscarmellose sodium or crospovidone), lubricants (e.g., silica, talc, stearic acid or salts thereof such as magnesium stearate, or polyethylene glycol), coating agents (e.g., concentrated sugar solutions including gum arabic, talc, polyvinyl pyrrolidone, carbopol gel, polyethylene glycol, or titanium dioxide), and antioxidants (e.g., sodium metabisulfite, sodium bisulfite, sodium sulfite, dextrose, phenols, and thiophenols).

In a preferred embodiment, a pharmaceutical composition of the invention comprises at least one nonaqueous, pharmaceutically acceptable solvent and an antitumor compound having a solubility in ethanol of at least about 100, 200, 300, 400, 500, 600, 700 or 800 mg/ml. While not being bound to a particular theory, it is believed that the ethanol solubility of the antitumor compound may be directly related to its efficacy. The antitumor compound can also be capable of being crystallized from a solution. In other words, a crystalline antitumor compound, such as compound 1393, can be dissolved in a solvent to form a solution and then recrystallized upon evaporation of the solvent without the formation of any



Express Mail No. EL 998648560 US

amorphous antitumor compound. It is also preferred that the antitumor compound have an ID<sub>50</sub> value (i.e, the drug concentration producing 50% inhibition of colony formation) of at least 4, 5, 6, 7, 8, 9, or 10 times less that of paclitaxel when measured according to the protocol set forth in the working examples.

Dosage form administration by these routes may be continuous or intermittent, depending, for example, upon the patient's physiological condition, whether the purpose of the administration is therapeutic or prophylactic, and other factors known to and assessable by a skilled practitioner.

Dosage and regimens for the administration of the pharmaceutical compositions of the invention can be readily determined by those with ordinary skill in treating cancer. It is understood that the dosage of the antitumor compounds will be dependent upon the age, sex, health, and weight of the recipient, kind of concurrent treatment, if any, frequency of treatment, and the nature of the effect desired. For any mode of administration, the actual amount of antitumor compound delivered, as well as the dosing schedule necessary to achieve the advantageous effects described herein, will also depend, in part, on such factors as the bioavailability of the antitumor compound, the disorder being treated, the desired therapeutic dose, and other factors that will be apparent to those of skill in the art. The dose administered to an animal, particularly a human, in the context of the present invention should be sufficient to effect the desired therapeutic response in the animal over a reasonable period of time. Preferably, an effective amount of the antitumor compound, whether administered orally or by another route, is any amount which would result in a desired therapeutic response when administered by that route. Preferably, the compositions for oral administration are prepared in such a way that a single dose in one or more oral preparations contains at least 20 mg of the antitumor compound per m<sup>2</sup> of patient body surface area, or at least 50, 100, 150, 200, 300, 400, or 500 mg of the antitumor compound per m<sup>2</sup> of patient body surface area, wherein the average body surface area for a human is 1.8 m<sup>2</sup>. Preferably, a single dose of a composition for oral administration contains from about 20 to about 600 mg of the antitumor compound per m<sup>2</sup> of patient body surface area, more preferably from about 25 to about 400 mg/m<sup>2</sup>, even more preferably, from about 40 to about 300 mg/m<sup>2</sup>, and even more preferably from about 50 to about 200 mg/m<sup>2</sup>. Preferably, the compositions for parenteral administration are

Express Mail No. EL 998648560 US

prepared in such a way that a single dose contains at least 20 mg of the antitumor compound per m<sup>2</sup> of patient body surface area, or at least 40, 50, 100, 150, 200, 300, 400, or 500 mg of the antitumor compound per m<sup>2</sup> of patient body surface area. Preferably, a single dose in one or more parenteral preparations contains from about 20 to about 500 mg of the antitumor compound per m<sup>2</sup> of patient body surface area, more preferably from about 40 to about 400 mg/m<sup>2</sup> and even more preferably, from about 60 to about 350 mg/m<sup>2</sup>. However, the dosage may vary depending on the dosing schedule which can be adjusted as necessary to achieve the desired therapeutic effect. It should be noted that the ranges of effective doses provided herein are not intended to limit the invention and represent preferred dose ranges. The most preferred dosage will be tailored to the individual subject, as is understood and determinable by one of ordinary skill in the art without undue experimentation.

The concentration of the antitumor compound in a liquid pharmaceutical composition is preferably between about 0.01 mg and about 10 mg per ml of the composition, more preferably between about 0.1 mg and about 7 mg per ml, even more preferably between about 0.5 mg and about 5 mg per ml, and most preferably between about 1.5 mg and about 4 mg per ml. Relatively low concentrations are generally preferred because the antitumor compound is most soluble in the solution at low concentrations. The concentration of the antitumor compound in a solid pharmaceutical composition for oral administration is preferably between about 5 weight % and about 50 weight %, based on the total weight of the composition, more preferably between about 8 weight % and about 40 weight %, and most preferably between about 10 weight % and about 30 weight %.

In one embodiment, solutions for oral administration are prepared by dissolving an antitumor compound in any pharmaceutically acceptable solvent capable of dissolving the compound (e.g., ethanol or methylene chloride) to form a solution. An appropriate volume of a carrier which is a solution, such as Cremophor® EL solution, is added to the solution while stirring to form a pharmaceutically acceptable solution for oral administration to a patient. If desired, such solutions can be formulated to contain a minimal amount of, or to be free of, ethanol, which is known in the art to cause adverse physiological effects when administered at certain concentrations in oral formulations.

Express Mail No. EL 998648560 US

In another embodiment, powders or tablets for oral administration are prepared by dissolving an antitumor compound in any pharmaceutically acceptable solvent capable of dissolving the compound (e.g., ethanol or methylene chloride) to form a solution. The solvent can optionally be capable of evaporating when the solution is dried under vacuum. An additional carrier can be added to the solution prior to drying, such as Cremophor® EL solution. The resulting solution is dried under vacuum to form a glass. The glass is then mixed with a binder to form a powder. The powder can be mixed with fillers or other conventional tableting agents and processed to form a tablet for oral administration to a patient. The powder can also be added to any liquid carrier as described above to form a solution, emulsion, suspension or the like for oral administration.

Emulsions for parenteral administration can be prepared by dissolving an antitumor compound in any pharmaceutically acceptable solvent capable of dissolving the compound (e.g., ethanol or methylene chloride) to form a solution. An appropriate volume of a carrier which is an emulsion, such as Liposyn® II or Liposyn® III emulsion, is added to the solution while stirring to form a pharmaceutically acceptable emulsion for parenteral administration to a patient. If desired, such emulsions can be formulated to contain a minimal amount of, or to be free of, ethanol or Cremophor® solution, which are known in the art to cause adverse physiological effects when administered at certain concentrations in parenteral formulations.

Solutions for parenteral administration can be prepared by dissolving an antitumor compound in any pharmaceutically acceptable solvent capable of dissolving the compound (e.g., ethanol or methylene chloride) to form a solution. An appropriate volume of a carrier which is a solution, such as Cremophor® solution, is added to the solution while stirring to form a pharmaceutically acceptable solution for parenteral administration to a patient. If desired, such solutions can be formulated to contain a minimal amount of, or to be free of, ethanol or Cremophor® solution, which are known in the art to cause adverse physiological effects when administered at certain concentrations in parenteral formulations.

If desired, the emulsions or solutions described above for oral or parenteral administration can be packaged in IV bags, vials or other conventional containers in concentrated form and diluted with any pharmaceutically acceptable liquid,

Express Mail No. EL 998648560 US

such as saline, to form an acceptable taxane concentration prior to use as is known in the art.

### Definitions

5 The terms "hydrocarbon" and "hydrocarbyl" as used herein describe organic compounds or radicals consisting exclusively of the elements carbon and hydrogen. These moieties include alkyl, alkenyl, alkynyl, and aryl moieties. These moieties also include alkyl, alkenyl, alkynyl, and aryl moieties substituted with other aliphatic or cyclic hydrocarbon groups, such as alkaryl, alkenaryl and alkynaryl. Unless otherwise indicated, these moieties preferably comprise 1 to 20  
10 carbon atoms.

The "substituted hydrocarbyl" moieties described herein are hydrocarbyl moieties which are substituted with at least one atom other than carbon, including moieties in which a carbon chain atom is substituted with a hetero atom such as nitrogen, oxygen, silicon, phosphorous, boron, sulfur, or a halogen atom. These  
15 substituents include halogen, heterocyclo, alkoxy, alkenoxy, alkynoxy, aryloxy, hydroxy, protected hydroxy, keto, acyl, acyloxy, nitro, amino, amido, nitro, cyano, thiol, ketals, acetals, esters and ethers.

The term "heteroatom" shall mean atoms other than carbon and hydrogen.

20 The "heterosubstituted methyl" moieties described herein are methyl groups in which the carbon atom is covalently bonded to at least one heteroatom and optionally with hydrogen, the heteroatom being, for example, a nitrogen, oxygen, silicon, phosphorous, boron, sulfur, or halogen atom. The heteroatom may, in turn, be substituted with other atoms to form a heterocyclo, alkoxy, alkenoxy, alkynoxy, aryloxy, hydroxy, protected hydroxy, oxy, acyloxy, nitro,  
25 amino, amido, thiol, ketals, acetals, esters or ether moiety.

30 The "heterosubstituted acetate" moieties described herein are acetate groups in which the carbon of the methyl group is covalently bonded to at least one heteroatom and optionally with hydrogen, the heteroatom being, for example, a nitrogen, oxygen, silicon, phosphorous, boron, sulfur, or halogen atom. The heteroatom may, in turn, be substituted with other atoms to form a heterocyclo, alkoxy, alkenoxy, alkynoxy, aryloxy, hydroxy, protected hydroxy, oxy, acyloxy, nitro, amino, amido, thiol, ketals, acetals, esters or ether moiety.

Unless otherwise indicated, the alkyl groups described herein are preferably lower alkyl containing from one to eight carbon atoms in the principal

Express Mail No. EL 998648560 US

chain and up to 20 carbon atoms. They may be straight or branched chain or cyclic and include methyl, ethyl, propyl, isopropyl, butyl, hexyl and the like.

Unless otherwise indicated, the alkenyl groups described herein are preferably lower alkenyl containing from two to eight carbon atoms in the principal chain and up to 20 carbon atoms. They may be straight or branched chain or cyclic and include ethenyl, propenyl, isopropenyl, butenyl, isobutenyl, hexenyl, and the like.

Unless otherwise indicated, the alkynyl groups described herein are preferably lower alkynyl containing from two to eight carbon atoms in the principal chain and up to 20 carbon atoms. They may be straight or branched chain and include ethynyl, propynyl, butynyl, isobutynyl, hexynyl, and the like.

The terms "aryl" or "ar" as used herein alone or as part of another group denote optionally substituted homocyclic aromatic groups, preferably monocyclic or bicyclic groups containing from 6 to 12 carbons in the ring portion, such as phenyl, biphenyl, naphthyl, substituted phenyl, substituted biphenyl or substituted naphthyl. Phenyl and substituted phenyl are the more preferred aryl.

The terms "halogen" or "halo" as used herein alone or as part of another group refer to chlorine, bromine, fluorine, and iodine.

The terms "heterocyclo" or "heterocyclic" as used herein alone or as part of another group denote optionally substituted, fully saturated or unsaturated, monocyclic or bicyclic, aromatic or nonaromatic groups having at least one heteroatom in at least one ring, and preferably 5 or 6 atoms in each ring. The heterocyclo group preferably has 1 or 2 oxygen atoms, 1 or 2 sulfur atoms, and/or 1 to 4 nitrogen atoms in the ring, and may be bonded to the remainder of the molecule through a carbon or heteroatom. Exemplary heterocyclo include heteroaromatics such as furyl, thienyl, pyridyl, oxazolyl, pyrrolyl, indolyl, quinoliny, or isoquinoliny and the like. Exemplary substituents include one or more of the following groups: hydrocarbyl, substituted hydrocarbyl, keto, hydroxy, protected hydroxy, acyl, acyloxy, alkoxy, alkenoxy, alkynoxy, aryloxy, halogen, amido, amino, nitro, cyano, thiol, ketals, acetals, esters and ethers.

The term "heteroaromatic" as used herein alone or as part of another group denote optionally substituted aromatic groups having at least one heteroatom in at least one ring, and preferably 5 or 6 atoms in each ring. The heteroaromatic group preferably has 1 or 2 oxygen atoms, 1 or 2 sulfur atoms, and/or 1 to 4 nitrogen atoms in the ring, and may be bonded to the remainder of

Express Mail No. EL 998648560 US

the molecule through a carbon or heteroatom. Exemplary heteroaromatics include furyl, thienyl, pyridyl, oxazolyl, pyrrolyl, indolyl, quinoliny, or isoquinoliny and the like. Exemplary substituents include one or more of the following groups: hydrocarbyl, substituted hydrocarbyl, keto, hydroxy, protected hydroxy, acyl, acyloxy, alkoxy, alkenoxy, alkynoxy, aryloxy, halogen, amido, amino, nitro, cyano, thiol, ketals, acetals, esters and ethers.

The term "acyl," as used herein alone or as part of another group, denotes the moiety formed by removal of the hydroxyl group from the group --COOH of an organic carboxylic acid, e.g., RC(O)-, wherein R is R<sup>1</sup>, R<sup>1</sup>O-, R<sup>1</sup>R<sup>2</sup>N-, or R<sup>1</sup>S-, R<sup>1</sup> is hydrocarbyl, heterosubstituted hydrocarbyl, or heterocyclo and R<sup>2</sup> is hydrogen, hydrocarbyl or substituted hydrocarbyl.

The term "acyloxy," as used herein alone or as part of another group, denotes an acyl group as described above bonded through an oxygen linkage (--O--), e.g., RC(O)O- wherein R is as defined in connection with the term "acyl."

Unless otherwise indicated, the alkoxycarbonyloxy moieties described herein comprise lower hydrocarbon or substituted hydrocarbon or substituted hydrocarbon moieties.

Unless otherwise indicated, the carbamoyloxy moieties described herein are derivatives of carbamic acid in which one or both of the amine hydrogens is optionally replaced by a hydrocarbyl, substituted hydrocarbyl or heterocyclo moiety.

The terms "hydroxyl protecting group" and "hydroxy protecting group" as used herein denote a group capable of protecting a free hydroxyl group ("protected hydroxyl") which, subsequent to the reaction for which protection is employed, may be removed without disturbing the remainder of the molecule. A variety of protecting groups for the hydroxyl group and the synthesis thereof may be found in "Protective Groups in Organic Synthesis" by T. W. Greene, John Wiley and Sons, 1981, or Fieser & Fieser. Exemplary hydroxyl protecting groups include methoxymethyl, 1-ethoxyethyl, benzyloxymethyl, (.beta.-trimethylsilylethoxy)methyl, tetrahydropyranyl, 2,2,2-trichloroethoxycarbonyl, t-butyl(diphenyl)silyl, trialkylsilyl, trichloromethoxycarbonyl and 2,2,2-trichloroethoxymethyl.


As used herein, "Ac" means acetyl; "Bz" means benzoyl; "Et" means ethyl; "Me" means methyl; "Ph" means phenyl; "iPr" means isopropyl; "tBu" and "t-Bu" means tert-butyl; "R" means lower alkyl unless otherwise defined; "py" means

Express Mail No. EL 998648560 US


pyridine or pyridyl; "TES" means triethylsilyl; "TMS" means trimethylsilyl; "LAH" means lithium aluminum hydride; "10-DAB" means 10-desacetylbaecatin III"; "amine protecting group" includes, but is not limited to, carbamates, for example, 2,2,2-trichloroethylcarbamate or tertbutylcarbamate; "protected hydroxy" means -  
5 OP wherein P is a hydroxy protecting group; "tBuOCO" and "BOC" mean tert-butoxycarbonyl; "tAmOCO" means tert-amylloxycarbonyl; "PhCO" means phenylcarbonyl; "2-FuCO" means 2-furylcarbonyl; "2-ThCO" means 2-thienylcarbonyl; "2-PyCO" means 2-pyridylcarbonyl; "3-PyCO" means 3-pyridylcarbonyl; "4-PyCO" means 4-pyridylcarbonyl; "C<sub>4</sub>H<sub>7</sub>CO" means  
10 butenylcarbonyl; "EtOCO" means ethoxycarbonyl; "ibueCO" means isobutenylcarbonyl; "iBuCO" means isobutylcarbonyl; "iBuOCO" means isobutoxycarbonyl; "iPrOCO" means isopropylloxycarbonyl; "nPrOCO" means n-propylloxycarbonyl; "nPrCO" means n-propylcarbonyl; "tC<sub>3</sub>H<sub>5</sub>CO" means *trans*-propenyl carbonyl; "ibue" means isobutenyl; "THF" means tetrahydrofuran;  
15 "DMAP" means 4-dimethylamino pyridine; "LHMDS" means Lithium HexamethylDiSilazide.

The following examples illustrate the invention.

Express Mail No. EL 998648560 US

Example 1

**10-Propionyl-10-deacetyl baccatin III.** To a mixture of 0.2 g (0.367 mmol) of 10-deacetyl baccatin III and 0.272 g (1.10 mmol) of  $\text{CeCl}_3$  in 10 mL of THF at 25 °C was added 2.35 mL (18.36 mmol) of propionic anhydride. After 30 min the reaction mixture was diluted with 200 mL of EtOAc, then washed three times with 50 mL of saturated aqueous  $\text{NaHCO}_3$  solution and brine. The organic extract was dried over  $\text{Na}_2\text{SO}_4$  and concentrated *in vacuo*. The crude solid was purified by flash column chromatography on silica gel using 70% EtOAc/hexane as eluent to give 0.199 g (90%) of 10-propionyl-10-deacetyl baccatin III as a solid.




**7-Dimethylphenylsilyl-10-propionyl-10-deacetyl baccatin III.** To a solution of 0.200 g (0.333 mmol) of 10-propionyl-10-deacetyl baccatin III in 12 mL of THF at -10 °C under a nitrogen atmosphere was added dropwise 0.668 mL (4.00 mmol) of chlorodimethyl-phenylsilane and 2.48 mL (30.64 mmol) of pyridine. After 90 min the mixture was diluted with 100 mL of a 1:1 mixture of ethyl acetate and hexane. The mixture was washed with 20 mL of saturated aqueous sodium bicarbonate solution and the organic layer separated. The aqueous layer was extracted with 30 mL of a 1:1 mixture of ethyl acetate and hexane, and the combined organic extracts were washed with brine, dried over  $\text{Na}_2\text{SO}_4$ , and concentrated *in vacuo*. The crude solid was purified by flash column chromatography on silica gel using




Express Mail No. EL 998648560 US

50% EtOAc/hexane as eluent to give 0.242 g (99%) of 7-dimethylphenylsilyl-10-propionyl-10-deacetyl baccatin III as a solid.



**7-Dimethylphenylsilyl-2'-O-triethylsilyl-3'-desphenyl-3'-(2-thienyl)-10-propionyl-10-deacetyl taxol.** To a solution of 0.400 g (0.544 mmol) of 7-dimethylphenylsilyl-10-propionyl-10-deacetyl baccatin III in 5.5 mL of THF at -45 °C under a nitrogen atmosphere was added 0.681 mL (0.681 mmol) of a 1M solution of LHMDS in THF. After 1 h, a solution of 0.317 g (0.818 mmol) of *cis*-N-benzoyl-3-triethylsilyloxy-4-(2-thienyl) azetidin-2-one in 3 mL of THF was added slowly. The mixture was warmed to 0 °C and after 3 h 10 mL of saturated aqueous sodium bicarbonate solution was added and the mixture was extracted three times with 50 mL of ethyl acetate. The combined organic extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated *in vacuo*. The crude product was purified by flash column chromatography on silica gel using 40% EtOAc/hexane as eluent to give 0.531 g (87%) of 7-dimethylphenylsilyl-2'-O-triethylsilyl-3'-desphenyl-3'-(2-thienyl)-10-propionyl-10-deacetyl taxol as a solid.



**3'-Desphenyl-3'-(2-thienyl)-10-propionyl-10-deacetyl taxol.** To a solution of 0.521 g (0.464 mmol) of 7-dimethylphenylsilyl-2'-O-triethylsilyl-3'-desphenyl-3'-(2-thienyl)-10-propionyl-10-deacetyl taxol in 2 mL of CH<sub>3</sub>CN and 2 mL of pyridine at 0 °C was added 0.5 mL of a solution of 30% HF in H<sub>2</sub>O. After 3 h 20 mL of a

Express Mail No. EL 998648560 US

saturated aqueous sodium bicarbonate solution was added and the mixture was extracted three times with 50 mL of ethyl acetate. The combined organic extracts were washed with brine, dried over Na<sub>2</sub>SO<sub>4</sub>, and concentrated *in vacuo*. The crude product was purified by flash column chromatography on silica gel using 70% EtOAc/hexane as eluent to give 0.405 g (100%) of 3'-desphenyl-3'-(2-thienyl)-10-propionyl-10-deacetyl taxol as a solid. m.p. 154-155 °C; [α]<sub>D</sub><sup>25</sup> = -45.0 (c 0.1 in CHCl<sub>3</sub>); Anal. Calcd. for C<sub>46</sub>H<sub>51</sub>NO<sub>14</sub>S: C, 63.22; H, 5.88; Found: C, 62.94; H, 5.97.

**3'-Desphenyl-3'-(2-thienyl)-10-propionyl-10-deacetyl  
taxol <sup>1</sup>H NMR data (CDCl<sub>3</sub>)**


Proton	ppm	pattern	J (Hz)
2'	4.78	dd	H3'(2.1), 2'OH(4.1)
2'OH	3.51	d	H2'(4.1)
3'	6.07	dd	NH(8.6), H2'(2.1)
5'	7.04	dd	(3.5), (5.0)
1OH	1.68	s	
2	5.69	d	H3(7.0)
3	3.85	d	H2(7.0)
4Ac	2.42	s	
5	4.96	app d	
6a	2.45-2.60	app m	
6b	1.89	ddd	H7(10.9), H5(2.5), H6a(14.5)
7	4.42	ddd	7OH(4.2), H6a(6.8), H6b(10.8)
7OH	2.45-2.60	app m	
10	6.32	s	
13	6.27	app t	H14a,b(9.0)
14a	2.40-2.43	app m	
14b	2.34	dd	H14a(15.5), H13(9.0)
Me 16	1.16	s	
Me 17	1.25	app m	
Me18	1.84	s	
Me19	1.70	s	

Express Mail No. EL 998648560 US

20a	4.31	d	H20b(8.5)
20b	4.22	d	H20a(8.5)
o-benzoate	8.14-8.16	m	
o-benzamide	7.72-7.73	m	
NH	6.88	d	H3'(8.6)
CH <sub>3</sub> CH <sub>2</sub>	1.24	t	CH <sub>3</sub> CH <sub>2</sub> (7.0)
CH <sub>3</sub> CH <sub>2</sub>	2.45-2.60	app m	

Example 2

The procedures described in Example 1 were repeated, but other suitably protected  $\beta$ -lactams were substituted for the  $\beta$ -lactam of Example 1 to prepare the series of compounds having structural formula (13) and the combinations of substituents identified in the following table



(13)

Compound	X <sub>5</sub>	X <sub>3</sub>	R <sub>10</sub>
0499	tBuOCO-	isobutenyl	EtCOO-
0503	tBuOCO-	2-pyridyl	EtCOO-
0517	tBuOCO-	3-pyridyl	EtCOO-
0521	tBuOCO-	4-pyridyl	EtCOO-
0536	tBuOCO-	2-furyl	EtCOO-
0549	tBuOCO-	3-furyl	EtCOO-
0550	tBuOCO-	2-thienyl	EtCOO-
0562	tBuOCO-	3-thienyl	EtCOO-

Express Mail No. EL 998648560 US

	0578	tBuOCO-	cyclopropyl	EtCOO-
	0583	tBuOCO-	isopropyl	EtCOO-
	0596	tBuOCO-	cyclobutyl	EtCOO-
	0602	tBuOCO-	p-nitrophenyl	EtCOO-
5	0611	tBuOCO-	phenyl	EtCOO-
	0625	PhCO-	isobutenyl	EtCOO-
	0634	PhCO-	2-pyridyl	EtCOO-
	0647	PhCO-	3-pyridyl	EtCOO-
	0659	PhCO-	4-pyridyl	EtCOO-
10	0663	PhCO-	2-furyl	EtCOO-
	0670	PhCO-	3-furyl	EtCOO-
	0687	PhCO-	2-thienyl	EtCOO-
	0691	PhCO-	3-thienyl	EtCOO-
	0706	PhCO-	cyclopropyl	EtCOO-
15	0719	PhCO-	isopropyl	EtCOO-
	0720	PhCO-	cyclobutyl	EtCOO-
	0732	PhCO-	p-nitrophenyl	EtCOO-
	0748	PhCO-	phenyl	EtCOO-
	0838	tBuOCO-	isobutenyl	cproCOO-
20	0843	tBuOCO-	2-furyl	cproCOO-
	0854	tBuOCO-	2-thienyl	cproCOO-
	0860	tBuOCO-	cyclopropyl	cproCOO-
	0879	tBuOCO-	p-nitrophenyl	cproCOO-
	0882	tBuOCO-	phenyl	cproCOO-
25	0890	PhCO-	isobutenyl	cproCOO-
	0908	PhCO-	2-furyl	cproCOO-
	0919	PhCO-	2-thienyl	cproCOO-
	0923	PhCO-	cyclopropyl	cproCOO-
	0937	PhCO-	phenyl	cproCOO-
30	0947	tBuOCO-	isobutenyl	PrCOO-

Express Mail No. EL 998648560 US

5

10

15

20

25

30

0951	tBuOCO-	2-pyridyl	PrCOO-
0966	tBuOCO-	3-pyridyl	PrCOO-
0978	tBuOCO-	4-pyridyl	PrCOO-
0983	tBuOCO-	2-furyl	PrCOO-
0999	tBuOCO-	3-furyl	PrCOO-
1003	tBuOCO-	2-thienyl	PrCOO-
1011	tBuOCO-	3-thienyl	PrCOO-
1020	tBuOCO-	cyclopropyl	PrCOO-
1031	tBuOCO-	isopropyl	PrCOO-
1044	tBuOCO-	cyclobutyl	PrCOO-
1060	tBuOCO-	phenyl	PrCOO-
1879	tBuOCO-	isobutenyl	2-ThCOO-
1883	tBuOCO-	2-pyridyl	2-ThCOO-
1892	tBuOCO-	2-furyl	2-ThCOO-
1900	tBuOCO-	2-thienyl	2-ThCOO-
1911	tBuOCO-	p-nitrophenyl	2-ThCOO-
1923	tBuOCO-	3-furyl	2-ThCOO-
1939	tBuOCO-	3-thienyl	2-ThCOO-
1948	tBuOCO-	3-pyridyl	2-ThCOO-
1954	tBuOCO-	4-pyridyl	2-ThCOO-
1964	tBuOCO-	isopropyl	2-ThCOO-
1970	tBuOCO-	cyclobutyl	2-ThCOO-
1988	tBuOCO-	phenyl	2-ThCOO-
2101	tBuOCO-	isobutenyl	2-FuCOO-
2111	tBuOCO-	2-pyridyl	2-FuCOO-
2124	tBuOCO-	3-pyridyl	2-FuCOO-
2132	tBuOCO-	4-pyridyl	2-FuCOO-
2142	tBuOCO-	2-furyl	2-FuCOO-
2159	tBuOCO-	3-furyl	2-FuCOO-
2164	tBuOCO-	2-thienyl	2-FuCOO-

Express Mail No. EL 998648560 US

2173	tBuOCO-	3-thienyl	2-FuCOO-
2181	tBuOCO-	isopropyl	2-FuCOO-
2199	tBuOCO-	cyclobutyl	2-FuCOO-
2202	tBuOCO-	p-nitrophenyl	2-FuCOO-
2212	tBuOCO-	phenyl	2-FuCOO-
2226	tBuOCO-	isobutenyl	iPrCOO-
2238	tBuOCO-	2-pyridyl	iPrCOO-
2242	tBuOCO-	3-pyridyl	iPrCOO-
2255	tBuOCO-	4-pyridyl	iPrCOO-
2269	tBuOCO-	2-furyl	iPrCOO-
2273	tBuOCO-	3-furyl	iPrCOO-
2287	tBuOCO-	2-thienyl	iPrCOO-
2291	tBuOCO-	3-thienyl	iPrCOO-
2306	tBuOCO-	isopropyl	iPrCOO-
2319	tBuOCO-	cyclobutyl	iPrCOO-
2320	tBuOCO-	p-nitrophenyl	iprCOO-
2332	tBuOCO-	isobutenyl	tC <sub>3</sub> H <sub>5</sub> COO-
2348	tBuOCO-	2-pyridyl	tC <sub>3</sub> H <sub>5</sub> COO-
2353	tBuOCO-	3-pyridyl	tC <sub>3</sub> H <sub>5</sub> COO-
2366	tBuOCO-	4-pyridyl	tC <sub>3</sub> H <sub>5</sub> COO-
2379	tBuOCO-	2-furyl	tC <sub>3</sub> H <sub>5</sub> COO-
2380	tBuOCO-	3-furyl	tC <sub>3</sub> H <sub>5</sub> COO-
2392	tBuOCO-	2-thienyl	tC <sub>3</sub> H <sub>5</sub> COO-
2408	tBuOCO-	3-thienyl	tC <sub>3</sub> H <sub>5</sub> COO-
2413	tBuOCO-	isopropyl	tC <sub>3</sub> H <sub>5</sub> COO-
2424	tBuOCO-	cyclobutyl	tC <sub>3</sub> H <sub>5</sub> COO-
2439	tBuOCO-	p-nitrophenyl	tC <sub>3</sub> H <sub>5</sub> COO-
2442	tBuOCO-	phenyl	tC <sub>3</sub> H <sub>5</sub> COO-
2455	tBuOCO-	isobutenyl	ibueCOO-
2464	tBuOCO-	2-pyridyl	ibueCOO-

Express Mail No. EL 998648560 US

2472	tBuOCO-	4-pyridyl	ibueCOO-
2488	tBuOCO-	2-furyl	ibueCOO-
2499	tBuOCO-	3-furyl	ibueCOO-
2503	tBuOCO-	2-thienyl	ibueCOO-
2511	tBuOCO-	3-thienyl	ibueCOO-
2520	tBuOCO-	phenyl	ibueCOO-
2781	tBuOCO-	3-furyl	cproCOO-
2794	tBuOCO-	3-thienyl	cproCOO-
2802	tBuOCO-	2-pyridyl	cproCOO-
2813	tBuOCO-	4-pyridyl	cproCOO-
2826	PhCO-	3-furyl	cproCOO-
2838	PhCO-	3-thienyl	cproCOO-
2844	PhCO-	2-pyridyl	cproCOO-
2855	PhCO-	4-pyridyl	cproCOO-
2869	PhCO-	p-nitrophenyl	cproCOO-
3053	2-FuCO-	2-thienyl	EtCOO-
3071	iPrOCO-	2-thienyl	cproCOO-
3096	EtOCO-	2-thienyl	PrCOO-
3102	iBuOCO-	2-furyl	cproCOO-
3110	iBuOCO-	2-furyl	PrCOO-
3129	iBuOCO-	2-thienyl	cproCOO-
3132	nPrCO-	2-thienyl	cproCOO-
3148	nPrCO-	2-thienyl	PrCOO-
3163	iBuOCO-	2-thienyl	EtCOO-
3204	PhCO-	2-furyl	PrCOO-
3219	nPrCO-	2-furyl	EtCOO-
3222	nPrCO-	2-furyl	PrCOO-
3258	PhCO-	2-thienyl	PrCOO-
3265	iBuOCO-	2-thienyl	PrCOO-
3297	2-FuCO-	2-thienyl	cproCOO-

Express Mail No. EL 998648560 US

	3314	nPrCO-	2-thienyl	PrCOO-
	3352	2-FuCO-	2-thienyl	PrCOO-
	3361	iPrOCO-	2-thienyl	EtCOO-
	3370	EtOCO-	2-thienyl	EtCOO-
5	3408	2-ThCO-	2-thienyl	PrCOO-
	3417	iPrOCO-	2-furyl	PrCOO-
	3425	2-ThCO-	2-thienyl	EtCOO-
	3453	2-ThCO-	2-thienyl	cproCOO-
	3482	PhCO-	cyclopropyl	PrCOO-
10	3494	tC <sub>3</sub> H <sub>5</sub> CO-	2-thienyl	EtCOO-
	3513	tC <sub>3</sub> H <sub>5</sub> CO-	2-thienyl	cproCOO-
	3522	iPrOCO-	2-furyl	EtCOO-
	3535	EtOCO-	2-furyl	EtCOO-
	3543	C <sub>4</sub> H <sub>7</sub> CO-	2-thienyl	cproCOO-
15	3588	C <sub>4</sub> H <sub>7</sub> CO-	2-thienyl	EtCOO-
	3595	tC <sub>3</sub> H <sub>5</sub> CO-	2-thienyl	PrCOO-
	3603	C <sub>4</sub> H <sub>7</sub> CO-	2-thienyl	PrCOO-
	3644	2-ThCO-	2-furyl	EtCOO-
	3656	2-ThCO-	2-furyl	PrCOO-
20	3663	2-ThCO-	2-furyl	cproCOO-
	3677	EtOCO-	2-furyl	cproCOO-
	3686	2-FuCO-	2-furyl	PrCOO-
	3693	EtOCO-	2-furyl	PrCOO-
	3800	C <sub>4</sub> H <sub>7</sub> CO-	2-furyl	PrCOO-
25	3818	2-FuCO-	2-furyl	EtCOO-
	3853	iPrOCO-	2-furyl	cproCOO-
	3866	2-FuCO-	2-furyl	cproCOO-
	3909	iPrOCO-	2-thienyl	PrCOO-
	3938	C <sub>4</sub> H <sub>7</sub> CO-	2-furyl	cproCOO-
30	3945	C <sub>4</sub> H <sub>7</sub> CO-	2-furyl	EtCOO-



Express Mail No. EL 998648560 US


	3957	iBuOCO-	2-furyl	PrCOO-
	3971	tC <sub>3</sub> H <sub>5</sub> CO-	2-furyl	cproCOO-
	3982	tC <sub>3</sub> H <sub>5</sub> CO-	2-furyl	EtCOO-
	3994	tC <sub>3</sub> H <sub>5</sub> CO-	2-furyl	PrCOO-
5	4051	EtOCO-	2-thienyl	cproCOO-
	4062	nPrCO-	2-furyl	cproCOO-
	4112	3-PyCO-	2-thienyl	cproCOO-
	4121	3-PyCO-	2-thienyl	EtCOO-
	4190	3-PyCO-	2-thienyl	PrCOO-
10	4207	4-PyCO-	2-thienyl	EtCOO-
	4329	ibueCO-	2-thienyl	cproCOO-
	4335	ibueCO-	2-thienyl	EtCOO-
	4344	ibueCO-	2-thienyl	PrCOO-
	4665	iBuOCO-	3-furyl	cproCOO-
15	4704	iBuOCO-	3-furyl	PrCOO-
	4711	iBuOCO-	3-thienyl	EtCOO-
	4720	iBuOCO-	isobutenyl	cproCOO-
	4799	iBuOCO-	cyclopropyl	EtCOO-
	4808	iBuOCO-	cyclopropyl	nPrCOO-
20	4834	iBuOCO-	3-thienyl	nPrCOO-
	4888	tC <sub>3</sub> H <sub>5</sub> CO-	3-furyl	EtCOO-
	4919	tC <sub>3</sub> H <sub>5</sub> CO-	3-furyl	nPrCOO-
	4944	tC <sub>3</sub> H <sub>5</sub> CO-	3-furyl	cproCOO-
	5011	iBuOCO-	3-thienyl	cproCOO-
25	5040	tC <sub>3</sub> H <sub>5</sub> CO-	3-thienyl	cproCOO-
	5065	iBuOCO-	isobutenyl	EtCOO-
	5144	iBuOCO-	isobutenyl	nPrCOO-
	5232	iBuOCO-	cyclopropyl	cproCOO-
	5495	tBuOCO-	3-furyl	EtCOO-
30	6522	tAmOCO-	2-furyl	EtCOO-

Express Mail No. EL 998648560 US

Example 3

Following the processes described in Example 1 and elsewhere herein, the following specific taxanes having structural formula 14 may be prepared, wherein  $R_{10}$  is as previously defined, including wherein  $R_{10}$  is  $R_{10a}COO^-$  and  $R_{10a}$  is

(i) substituted or unsubstituted  $C_2$  to  $C_8$  alkyl such as ethyl, or straight, branched or cyclic propyl, butyl, pentyl, or hexyl; (ii) substituted or unsubstituted  $C_2$  to  $C_8$  alkenyl such as ethenyl or straight, branched or cyclic propenyl, butenyl, pentenyl or hexenyl; (iii) substituted or unsubstituted  $C_2$  to  $C_8$  alkynyl such as ethynyl or straight or branched propynyl, butynyl, pentynyl, or hexynyl; (iv) substituted or unsubstituted phenyl, or (v) substituted or unsubstituted heteroaromatic such as furyl, thienyl, or pyridyl. The substituents may be those identified elsewhere herein for substituted hydrocarbonyl. In one embodiment,  $R_{10}$  may be  $R_{10a}COO^-$  wherein  $R_{10a}$  is ethyl, straight, branched or cyclic propyl, straight or branched propenyl, isobutenyl, furyl or thienyl.



$X_5$	$X_3$	$R_{10}$
tBuOCO-	2-furyl	$R_aCOO^-$
tBuOCO-	3-furyl	$R_aCOO^-$
tBuOCO-	2-thienyl	$R_aCOO^-$
tBuOCO-	3-thienyl	$R_aCOO^-$
tBuOCO-	2-pyridyl	$R_aCOO^-$
tBuOCO-	3-pyridyl	$R_aCOO^-$
tBuOCO-	4-pyridyl	$R_aCOO^-$
tBuOCO-	isobutenyl	$R_aCOO^-$
tBuOCO-	isopropyl	$R_aCOO^-$

Express Mail No. EL 998648560 US

5

10

15

20

25

30

tBuOCO-	cyclopropyl	R <sub>a</sub> COO-
tBuOCO-	cyclobutyl	R <sub>a</sub> COO-
tBuOCO-	cyclopentyl	R <sub>a</sub> COO-
tBuOCO-	phenyl	R <sub>a</sub> COO-
benzoyl	2-furyl	R <sub>a</sub> COO-
benzoyl	3-furyl	R <sub>a</sub> COO-
benzoyl	2-thienyl	R <sub>a</sub> COO-
benzoyl	3-thienyl	R <sub>a</sub> COO-
benzoyl	2-pyridyl	R <sub>a</sub> COO-
benzoyl	3-pyridyl	R <sub>a</sub> COO-
benzoyl	4-pyridyl	R <sub>a</sub> COO-
benzoyl	isobutenyl	R <sub>a</sub> COO-
benzoyl	isopropyl	R <sub>a</sub> COO-
benzoyl	cyclopropyl	R <sub>a</sub> COO-
benzoyl	cyclobutyl	R <sub>a</sub> COO-
benzoyl	cyclopentyl	R <sub>a</sub> COO-
benzoyl	phenyl	R <sub>a</sub> COO-
2-FuCO-	2-furyl	R <sub>a</sub> COO-
2-FuCO-	3-furyl	R <sub>a</sub> COO-
2-FuCO-	2-thienyl	R <sub>a</sub> COO-
2-FuCO-	3-thienyl	R <sub>a</sub> COO-
2-FuCO-	2-pyridyl	R <sub>a</sub> COO-
2-FuCO-	3-pyridyl	R <sub>a</sub> COO-
2-FuCO-	4-pyridyl	R <sub>a</sub> COO-
2-FuCO-	isobutenyl	R <sub>a</sub> COO-
2-FuCO-	isopropyl	R <sub>a</sub> COO-
2-FuCO-	cyclopropyl	R <sub>a</sub> COO-
2-FuCO-	cyclobutyl	R <sub>a</sub> COO-
2-FuCO-	cyclopentyl	R <sub>a</sub> COO-
2-FuCO-	phenyl	R <sub>a</sub> COO-

Express Mail No. EL 998648560 US

5

10

15

20

25

30

2-ThCO-	2-furyl	R <sub>a</sub> COO-
2-ThCO-	3-furyl	R <sub>a</sub> COO-
2-ThCO-	2-thienyl	R <sub>a</sub> COO-
2-ThCO-	3-thienyl	R <sub>a</sub> COO-
2-ThCO-	2-pyridyl	R <sub>a</sub> COO-
2-ThCO-	3-pyridyl	R <sub>a</sub> COO-
2-ThCO-	4-pyridyl	R <sub>a</sub> COO-
2-ThCO-	isobutenyl	R <sub>a</sub> COO-
2-ThCO-	isopropyl	R <sub>a</sub> COO-
2-ThCO-	cyclopropyl	R <sub>a</sub> COO-
2-ThCO-	cyclobutyl	R <sub>a</sub> COO-
2-ThCO-	cyclopentyl	R <sub>a</sub> COO-
2-ThCO-	phenyl	R <sub>a</sub> COO-
2-PyCO-	2-furyl	R <sub>a</sub> COO-
2-PyCO-	3-furyl	R <sub>a</sub> COO-
2-PyCO-	2-thienyl	R <sub>a</sub> COO-
2-PyCO-	3-thienyl	R <sub>a</sub> COO-
2-PyCO-	2-pyridyl	R <sub>a</sub> COO-
2-PyCO-	3-pyridyl	R <sub>a</sub> COO-
2-PyCO-	4-pyridyl	R <sub>a</sub> COO-
2-PyCO-	isobutenyl	R <sub>a</sub> COO-
2-PyCO-	isopropyl	R <sub>a</sub> COO-
2-PyCO-	cyclopropyl	R <sub>a</sub> COO-
2-PyCO-	cyclobutyl	R <sub>a</sub> COO-
2-PyCO-	cyclopentyl	R <sub>a</sub> COO-
2-PyCO-	phenyl	R <sub>a</sub> COO-
3-PyCO-	2-furyl	R <sub>a</sub> COO-
3-PyCO-	3-furyl	R <sub>a</sub> COO-
3-PyCO-	2-thienyl	R <sub>a</sub> COO-
3-PyCO-	3-thienyl	R <sub>a</sub> COO-

Express Mail No. EL 998648560 US

5

10

15

20

25

30

3-PyCO-	2-pyridyl	R <sub>a</sub> COO-
3-PyCO-	3-pyridyl	R <sub>a</sub> COO-
3-PyCO-	4-pyridyl	R <sub>a</sub> COO-
3-PyCO-	isobutenyl	R <sub>a</sub> COO-
3-PyCO-	isopropyl	R <sub>a</sub> COO-
3-PyCO-	cyclopropyl	R <sub>a</sub> COO-
3-PyCO-	cyclobutyl	R <sub>a</sub> COO-
3-PyCO-	cyclopentyl	R <sub>a</sub> COO-
3-PyCO-	phenyl	R <sub>a</sub> COO-
4-PyCO-	2-furyl	R <sub>a</sub> COO-
4-PyCO-	3-furyl	R <sub>a</sub> COO-
4-PyCO-	2-thienyl	R <sub>a</sub> COO-
4-PyCO-	3-thienyl	R <sub>a</sub> COO-
4-PyCO-	2-pyridyl	R <sub>a</sub> COO-
4-PyCO-	3-pyridyl	R <sub>a</sub> COO-
4-PyCO-	4-pyridyl	R <sub>a</sub> COO-
4-PyCO-	isobutenyl	R <sub>a</sub> COO-
4-PyCO-	isopropyl	R <sub>a</sub> COO-
4-PyCO-	cyclopropyl	R <sub>a</sub> COO-
4-PyCO-	cyclobutyl	R <sub>a</sub> COO-
4-PyCO-	cyclopentyl	R <sub>a</sub> COO-
4-PyCO-	phenyl	R <sub>a</sub> COO-
C <sub>4</sub> H <sub>7</sub> CO-	2-furyl	R <sub>a</sub> COO-
C <sub>4</sub> H <sub>7</sub> CO-	3-furyl	R <sub>a</sub> COO-
C <sub>4</sub> H <sub>7</sub> CO-	2-thienyl	R <sub>a</sub> COO-
C <sub>4</sub> H <sub>7</sub> CO-	3-thienyl	R <sub>a</sub> COO-
C <sub>4</sub> H <sub>7</sub> CO-	2-pyridyl	R <sub>a</sub> COO-
C <sub>4</sub> H <sub>7</sub> CO-	3-pyridyl	R <sub>a</sub> COO-
C <sub>4</sub> H <sub>7</sub> CO-	4-pyridyl	R <sub>a</sub> COO-
C <sub>4</sub> H <sub>7</sub> CO-	isobutenyl	R <sub>a</sub> COO-

Express Mail No. EL 998648560 US

5

10

15

20

25

30

C <sub>4</sub> H <sub>7</sub> CO-	isopropyl	R <sub>a</sub> COO-
C <sub>4</sub> H <sub>7</sub> CO-	cyclopropyl	R <sub>a</sub> COO-
C <sub>4</sub> H <sub>7</sub> CO-	cyclobutyl	R <sub>a</sub> COO-
C <sub>4</sub> H <sub>7</sub> CO-	cyclopentyl	R <sub>a</sub> COO-
C <sub>4</sub> H <sub>7</sub> CO-	phenyl	R <sub>a</sub> COO-
EtOCO-	2-furyl	R <sub>a</sub> COO-
EtOCO-	3-furyl	R <sub>a</sub> COO-
EtOCO-	2-thienyl	R <sub>a</sub> COO-
EtOCO-	3-thienyl	R <sub>a</sub> COO-
EtOCO-	2-pyridyl	R <sub>a</sub> COO-
EtOCO-	3-pyridyl	R <sub>a</sub> COO-
EtOCO-	4-pyridyl	R <sub>a</sub> COO-
EtOCO-	isobutenyl	R <sub>a</sub> COO-
EtOCO-	isopropyl	R <sub>a</sub> COO-
EtOCO-	cyclopropyl	R <sub>a</sub> COO-
EtOCO-	cyclobutyl	R <sub>a</sub> COO-
EtOCO-	cyclopentyl	R <sub>a</sub> COO-
EtOCO-	phenyl	R <sub>a</sub> COO-
ibueCO-	2-furyl	R <sub>a</sub> COO-
ibueCO-	3-furyl	R <sub>a</sub> COO-
ibueCO-	2-thienyl	R <sub>a</sub> COO-
ibueCO-	3-thienyl	R <sub>a</sub> COO-
ibueCO-	2-pyridyl	R <sub>a</sub> COO-
ibueCO-	3-pyridyl	R <sub>a</sub> COO-
ibueCO-	4-pyridyl	R <sub>a</sub> COO-
ibueCO-	isobutenyl	R <sub>a</sub> COO-
ibueCO-	isopropyl	R <sub>a</sub> COO-
ibueCO-	cyclopropyl	R <sub>a</sub> COO-
ibueCO-	cyclobutyl	R <sub>a</sub> COO-
ibueCO-	cyclopentyl	R <sub>a</sub> COO-

Express Mail No. EL 998648560 US

	ibueCO-	phenyl	R <sub>a</sub> COO-
	iBuCO-	2-furyl	R <sub>a</sub> COO-
	iBuCO-	3-furyl	R <sub>a</sub> COO-
5	iBuCO-	2-thienyl	R <sub>a</sub> COO-
	iBuCO-	3-thienyl	R <sub>a</sub> COO-
	iBuCO-	2-pyridyl	R <sub>a</sub> COO-
	iBuCO-	3-pyridyl	R <sub>a</sub> COO-
	iBuCO-	4-pyridyl	R <sub>a</sub> COO-
10	iBuCO-	isobutenyl	R <sub>a</sub> COO-
	iBuCO-	isopropyl	R <sub>a</sub> COO-
	iBuCO-	cyclopropyl	R <sub>a</sub> COO-
	iBuCO-	cyclobutyl	R <sub>a</sub> COO-
	iBuCO-	cyclopentyl	R <sub>a</sub> COO-
15	iBuCO-	phenyl	R <sub>a</sub> COO-
	iBuOCO-	2-furyl	R <sub>a</sub> COO-
	iBuOCO-	3-furyl	R <sub>a</sub> COO-
	iBuOCO-	2-thienyl	R <sub>a</sub> COO-
	iBuOCO-	3-thienyl	R <sub>a</sub> COO-
20	iBuOCO-	2-pyridyl	R <sub>a</sub> COO-
	iBuOCO-	3-pyridyl	R <sub>a</sub> COO-
	iBuOCO-	4-pyridyl	R <sub>a</sub> COO-
	iBuOCO-	isobutenyl	R <sub>a</sub> COO-
	iBuOCO-	isopropyl	R <sub>a</sub> COO-
25	iBuOCO-	cyclopropyl	R <sub>a</sub> COO-
	iBuOCO-	cyclobutyl	R <sub>a</sub> COO-
	iBuOCO-	cyclopentyl	R <sub>a</sub> COO-
	iBuCO-	phenyl	R <sub>a</sub> COO-
	iPrOCO-	2-furyl	R <sub>a</sub> COO-
30	iPrOCO-	3-furyl	R <sub>a</sub> COO-
	iPrOCO-	2-thienyl	R <sub>a</sub> COO-

Express Mail No. EL 998648560 US

5

10

15

20

25

30

iPrOCO-	3-thienyl	R <sub>a</sub> COO-
iPrOCO-	2-pyridyl	R <sub>a</sub> COO-
iPrOCO-	3-pyridyl	R <sub>a</sub> COO-
iPrOCO-	4-pyridyl	R <sub>a</sub> COO-
iPrOCO-	isobutenyl	R <sub>a</sub> COO-
iPrOCO-	isopropyl	R <sub>a</sub> COO-
iPrOCO-	cyclopropyl	R <sub>a</sub> COO-
iPrOCO-	cyclobutyl	R <sub>a</sub> COO-
iPrOCO-	cyclopentyl	R <sub>a</sub> COO-
iPrOCO-	phenyl	R <sub>a</sub> COO-
nPrOCO-	2-furyl	R <sub>a</sub> COO-
nPrOCO-	3-furyl	R <sub>a</sub> COO-
nPrOCO-	2-thienyl	R <sub>a</sub> COO-
nPrOCO-	3-thienyl	R <sub>a</sub> COO-
nPrOCO-	2-pyridyl	R <sub>a</sub> COO-
nPrOCO-	3-pyridyl	R <sub>a</sub> COO-
nPrOCO-	4-pyridyl	R <sub>a</sub> COO-
nPrOCO-	isobutenyl	R <sub>a</sub> COO-
nPrOCO-	isopropyl	R <sub>a</sub> COO-
nPrOCO-	cyclopropyl	R <sub>a</sub> COO-
nPrOCO-	cyclobutyl	R <sub>a</sub> COO-
nPrOCO-	cyclopentyl	R <sub>a</sub> COO-
nPrOCO-	phenyl	R <sub>a</sub> COO-
nPrCO-	2-furyl	R <sub>a</sub> COO-
nPrCO-	3-furyl	R <sub>a</sub> COO-
nPrCO-	2-thienyl	R <sub>a</sub> COO-
nPrCO-	3-thienyl	R <sub>a</sub> COO-
nPrCO-	2-pyridyl	R <sub>a</sub> COO-
nPrCO-	3-pyridyl	R <sub>a</sub> COO-
nPrCO-	4-pyridyl	R <sub>a</sub> COO-



Express Mail No. EL 998648560 US

5

10

15

20

25

30

nPrCO-	isobutenyl	R <sub>a</sub> COO-
nPrCO-	isopropyl	R <sub>a</sub> COO-
nPrCO-	cyclopropyl	R <sub>a</sub> COO-
nPrCO-	cyclobutyl	R <sub>a</sub> COO-
nPrCO-	cyclopentyl	R <sub>a</sub> COO-
nPrOCO-	phenyl	R <sub>a</sub> COO-
tBuOCO-	cyclopentyl	EtCOO-
benzoyl	cyclopentyl	EtCOO-
2-FuCO-	3-furyl	EtCOO-
2-FuCO-	3-thienyl	EtCOO-
2-FuCO-	2-pyridyl	EtCOO-
2-FuCO-	3-pyridyl	EtCOO-
2-FuCO-	4-pyridyl	EtCOO-
2-FuCO-	isobutenyl	EtCOO-
2-FuCO-	isopropyl	EtCOO-
2-FuCO-	cyclopropyl	EtCOO-
2-FuCO-	cyclobutyl	EtCOO-
2-FuCO-	cyclopentyl	EtCOO-
2-FuCO-	phenyl	EtCOO-
2-ThCO-	3-furyl	EtCOO-
2-ThCO-	3-thienyl	EtCOO-
2-ThCO-	2-pyridyl	EtCOO-
2-ThCO-	3-pyridyl	EtCOO-
2-ThCO-	4-pyridyl	EtCOO-
2-ThCO-	isobutenyl	EtCOO-
2-ThCO-	isopropyl	EtCOO-
2-ThCO-	cyclopropyl	EtCOO-
2-ThCO-	cyclobutyl	EtCOO-
2-ThCO-	cyclopentyl	EtCOO-
2-ThCO-	phenyl	EtCOO-

Express Mail No. EL 998648560 US

5

10

15

20

25

30

2-PyCO-	2-furyl	EtCOO-
2-PyCO-	3-furyl	EtCOO-
2-PyCO-	2-thienyl	EtCOO-
2-PyCO-	3-thienyl	EtCOO-
2-PyCO-	2-pyridyl	EtCOO-
2-PyCO-	3-pyridyl	EtCOO-
2-PyCO-	4-pyridyl	EtCOO-
2-PyCO-	isobutenyl	EtCOO-
2-PyCO-	isopropyl	EtCOO-
2-PyCO-	cyclopropyl	EtCOO-
2-PyCO-	cyclobutyl	EtCOO-
2-PyCO-	cyclopentyl	EtCOO-
2-PyCO-	phenyl	EtCOO-
3-PyCO-	2-furyl	EtCOO-
3-PyCO-	3-furyl	EtCOO-
3-PyCO-	3-thienyl	EtCOO-
3-PyCO-	2-pyridyl	EtCOO-
3-PyCO-	3-pyridyl	EtCOO-
3-PyCO-	4-pyridyl	EtCOO-
3-PyCO-	isobutenyl	EtCOO-
3-PyCO-	isopropyl	EtCOO-
3-PyCO-	cyclopropyl	EtCOO-
3-PyCO-	cyclobutyl	EtCOO-
3-PyCO-	cyclopentyl	EtCOO-
3-PyCO-	phenyl	EtCOO-
4-PyCO-	2-furyl	EtCOO-
4-PyCO-	3-furyl	EtCOO-
4-PyCO-	3-thienyl	EtCOO-
4-PyCO-	2-pyridyl	EtCOO-
4-PyCO-	3-pyridyl	EtCOO-

Express Mail No. EL 998648560 US

5

10

15

20

25

30

4-PyCO-	4-pyridyl	EtCOO-
4-PyCO-	isobutenyl	EtCOO-
4-PyCO-	isopropyl	EtCOO-
4-PyCO-	cyclopropyl	EtCOO-
4-PyCO-	cyclobutyl	EtCOO-
4-PyCO-	cyclopentyl	EtCOO-
4-PyCO-	phenyl	EtCOO-
C <sub>4</sub> H <sub>7</sub> CO-	3-furyl	EtCOO-
C <sub>4</sub> H <sub>7</sub> CO-	3-thienyl	EtCOO-
C <sub>4</sub> H <sub>7</sub> CO-	2-pyridyl	EtCOO-
C <sub>4</sub> H <sub>7</sub> CO-	3-pyridyl	EtCOO-
C <sub>4</sub> H <sub>7</sub> CO-	4-pyridyl	EtCOO-
C <sub>4</sub> H <sub>7</sub> CO-	isobutenyl	EtCOO-
C <sub>4</sub> H <sub>7</sub> CO-	isopropyl	EtCOO-
C <sub>4</sub> H <sub>7</sub> CO-	cyclopropyl	EtCOO-
C <sub>4</sub> H <sub>7</sub> CO-	cyclobutyl	EtCOO-
C <sub>4</sub> H <sub>7</sub> CO-	cyclopentyl	EtCOO-
C <sub>4</sub> H <sub>7</sub> CO-	phenyl	EtCOO-
EtOCO-	3-furyl	EtCOO-
EtOCO-	3-thienyl	EtCOO-
EtOCO-	2-pyridyl	EtCOO-
EtOCO-	3-pyridyl	EtCOO-
EtOCO-	4-pyridyl	EtCOO-
EtOCO-	isobutenyl	EtCOO-
EtOCO-	isopropyl	EtCOO-
EtOCO-	cyclopropyl	EtCOO-
EtOCO-	cyclobutyl	EtCOO-
EtOCO-	cyclopentyl	EtCOO-
EtOCO-	phenyl	EtCOO-
ibueCO-	2-furyl	EtCOO-

Express Mail No. EL 998648560 US

5

10

15

20

25

30

ibueCO-	3-furyl	EtCOO-
ibueCO-	3-thienyl	EtCOO-
ibueCO-	2-pyridyl	EtCOO-
ibueCO-	3-pyridyl	EtCOO-
ibueCO-	4-pyridyl	EtCOO-
ibueCO-	isobutenyl	EtCOO-
ibueCO-	isopropyl	EtCOO-
ibueCO-	cyclopropyl	EtCOO-
ibueCO-	cyclobutyl	EtCOO-
ibueCO-	cyclopentyl	EtCOO-
ibueCO-	phenyl	EtCOO-
iBuCO-	2-furyl	EtCOO-
iBuCO-	3-furyl	EtCOO-
iBuCO-	2-thienyl	EtCOO-
iBuCO-	3-thienyl	EtCOO-
iBuCO-	2-pyridyl	EtCOO-
iBuCO-	3-pyridyl	EtCOO-
iBuCO-	4-pyridyl	EtCOO-
iBuCO-	isobutenyl	EtCOO-
iBuCO-	isopropyl	EtCOO-
iBuCO-	cyclopropyl	EtCOO-
iBuCO-	cyclobutyl	EtCOO-
iBuCO-	cyclopentyl	EtCOO-
iBuCO-	phenyl	EtCOO-
iBuOCO-	2-furyl	EtCOO-
iBuOCO-	2-pyridyl	EtCOO-
iBuOCO-	3-pyridyl	EtCOO-
iBuOCO-	4-pyridyl	EtCOO-
iBuOCO-	isopropyl	EtCOO-
iBuOCO-	cyclobutyl	EtCOO-

Express Mail No. EL 998648560 US

5

10

15

20

25

30

iBuOCO-	cyclopentyl	EtCOO-
iBuCO-	phenyl	EtCOO-
iPrOCO-	3-furyl	EtCOO-
iPrOCO-	3-thienyl	EtCOO-
iPrOCO-	2-pyridyl	EtCOO-
iPrOCO-	3-pyridyl	EtCOO-
iPrOCO-	4-pyridyl	EtCOO-
iPrOCO-	isobutenyl	EtCOO-
iPrOCO-	isopropyl	EtCOO-
iPrOCO-	cyclopropyl	EtCOO-
iPrOCO-	cyclobutyl	EtCOO-
iPrOCO-	cyclopentyl	EtCOO-
iPrOCO-	phenyl	EtCOO-
nPrOCO-	2-furyl	EtCOO-
nPrOCO-	3-furyl	EtCOO-
nPrOCO-	2-thienyl	EtCOO-
nPrOCO-	3-thienyl	EtCOO-
nPrOCO-	2-pyridyl	EtCOO-
nPrOCO-	3-pyridyl	EtCOO-
nPrOCO-	4-pyridyl	EtCOO-
nPrOCO-	isobutenyl	EtCOO-
nPrOCO-	isopropyl	EtCOO-
nPrOCO-	cyclopropyl	EtCOO-
nPrOCO-	cyclobutyl	EtCOO-
nPrOCO-	cyclopentyl	EtCOO-
nPrOCO-	phenyl	EtCOO-
nPrCO-	3-furyl	EtCOO-
nPrCO-	3-thienyl	EtCOO-
nPrCO-	2-pyridyl	EtCOO-
nPrCO-	3-pyridyl	EtCOO-

Express Mail No. EL 998648560 US

nPrCO-	4-pyridyl	EtCOO-
nPrCO-	isobutenyl	EtCOO-
nPrCO-	isopropyl	EtCOO-
nPrCO-	cyclopropyl	EtCOO-
nPrCO-	cyclobutyl	EtCOO-
nPrCO-	cyclopentyl	EtCOO-
nPrOCO-	phenyl	EtCOO-

Example 4

Following the processes described in Example 1 and elsewhere herein, the following specific taxanes having structural formula 15 may be prepared, wherein  $R_7$  is hydroxy and  $R_{10}$  in each of the series (that is, each of series "A" through "K") is as previously defined, including wherein  $R_{10}$  is  $R_{10a}COO-$  and  $R_{10a}$  is (i) substituted or unsubstituted, preferably unsubstituted,  $C_2$  to  $C_8$  alkyl (straight, branched or cyclic), such as ethyl, propyl, butyl, pentyl, or hexyl; (ii) substituted or unsubstituted, preferably unsubstituted,  $C_2$  to  $C_8$  alkenyl (straight, branched or cyclic), such as ethenyl, propenyl, butenyl, pentenyl or hexenyl; (iii) substituted or unsubstituted, preferably unsubstituted,  $C_2$  to  $C_8$  alkynyl (straight or branched) such as ethynyl, propynyl, butynyl, pentynyl, or hexynyl; (iv) substituted or unsubstituted, preferably unsubstituted, phenyl; or (v) substituted or unsubstituted, preferably unsubstituted, heteroaromatic such as furyl, thienyl, or pyridyl.

Express Mail No. EL 998648560 US

In the "A" series of compounds,  $X_{10}$  is as otherwise as defined herein. Preferably, heterocyclo is substituted or unsubstituted furyl, thienyl, or pyridyl,  $X_{10}$  is substituted or unsubstituted furyl, thienyl, pyridyl, phenyl, or lower alkyl (e.g., tert-butyl), and  $R_7$  and  $R_{10}$  each have the beta stereochemical configuration.

5 In the "B" series of compounds,  $X_{10}$  and  $R_{2a}$  are as otherwise as defined herein. Preferably, heterocyclo is preferably substituted or unsubstituted furyl, thienyl, or pyridyl,  $X_{10}$  is preferably substituted or unsubstituted furyl, thienyl, pyridyl, phenyl, or lower alkyl (e.g., tert-butyl),  $R_{2a}$  is preferably substituted or unsubstituted furyl, thienyl, pyridyl, phenyl, or lower alkyl, and  $R_7$  and  $R_{10}$  each have the beta stereochemical configuration.

10 In the "C" series of compounds,  $X_{10}$  and  $R_{9a}$  are as otherwise as defined herein. Preferably, heterocyclo is preferably substituted or unsubstituted furyl, thienyl, or pyridyl,  $X_{10}$  is preferably substituted or unsubstituted furyl, thienyl, pyridyl, phenyl, or lower alkyl (e.g., tert-butyl),  $R_{9a}$  is preferably substituted or unsubstituted furyl, thienyl, pyridyl, phenyl, or lower alkyl, and  $R_7$ ,  $R_9$  and  $R_{10}$  each have the beta stereochemical configuration.

15 In the "D" and "E" series of compounds,  $X_{10}$  is as otherwise as defined herein. Preferably, heterocyclo is preferably substituted or unsubstituted furyl, thienyl, or pyridyl,  $X_{10}$  is preferably substituted or unsubstituted furyl, thienyl, pyridyl, phenyl, or lower alkyl (e.g., tert-butyl), and  $R_7$ ,  $R_9$  (series D only) and  $R_{10}$  each have the beta stereochemical configuration.

20 In the "F" series of compounds,  $X_{10}$ ,  $R_{2a}$  and  $R_{9a}$  are as otherwise as defined herein. Preferably, heterocyclo is preferably substituted or unsubstituted furyl, thienyl, or pyridyl,  $X_{10}$  is preferably substituted or unsubstituted furyl, thienyl, pyridyl, phenyl, or lower alkyl (e.g., tert-butyl),  $R_{2a}$  is preferably substituted or unsubstituted furyl, thienyl, pyridyl, phenyl, or lower alkyl, and  $R_7$ ,  $R_9$  and  $R_{10}$  each have the beta stereochemical configuration.

25 In the "G" series of compounds,  $X_{10}$  and  $R_{2a}$  are as otherwise as defined herein. Preferably, heterocyclo is preferably substituted or unsubstituted furyl, thienyl, or pyridyl,  $X_{10}$  is preferably substituted or unsubstituted furyl, thienyl, pyridyl, phenyl, or lower alkyl (e.g., tert-butyl),  $R_{2a}$  is preferably substituted or unsubstituted furyl, thienyl, pyridyl, phenyl, or lower alkyl, and  $R_7$ ,  $R_9$  and  $R_{10}$  each have the beta stereochemical configuration.

30 In the "H" series of compounds,  $X_{10}$  is as otherwise as defined herein. Preferably, heterocyclo is preferably substituted or unsubstituted furyl, thienyl, or

35

Express Mail No. EL 998648560 US


pyridyl,  $X_{10}$  is preferably substituted or unsubstituted furyl, thienyl, pyridyl, phenyl, or lower alkyl (e.g., tert-butyl),  $R_{2a}$  is preferably substituted or unsubstituted furyl, thienyl, pyridyl, phenyl, or lower alkyl, and  $R_7$  and  $R_{10}$  each have the beta stereochemical configuration.

5 In the "I" series of compounds,  $X_{10}$  and  $R_{2a}$  are as otherwise as defined herein. Preferably, heterocyclo is preferably substituted or unsubstituted furyl, thienyl, or pyridyl,  $X_{10}$  is preferably substituted or unsubstituted furyl, thienyl, pyridyl, phenyl, or lower alkyl (e.g., tert-butyl),  $R_{2a}$  is preferably substituted or unsubstituted furyl, thienyl, pyridyl, phenyl, or lower alkyl, and  $R_7$  and  $R_{10}$  each have the beta stereochemical configuration.

10 In the "J" series of compounds,  $X_{10}$  and  $R_{2a}$  are as otherwise as defined herein. Preferably, heterocyclo is preferably substituted or unsubstituted furyl, thienyl, or pyridyl,  $X_{10}$  is preferably substituted or unsubstituted furyl, thienyl, pyridyl, phenyl, or lower alkyl (e.g., tert-butyl),  $R_{2a}$  is preferably substituted or unsubstituted furyl, thienyl, pyridyl, phenyl, or lower alkyl, and  $R_7$ ,  $R_9$  and  $R_{10}$  each have the beta stereochemical configuration.

15 In the "K" series of compounds,  $X_{10}$ ,  $R_{2a}$  and  $R_{9a}$  are as otherwise as defined herein. Preferably, heterocyclo is preferably substituted or unsubstituted furyl, thienyl, or pyridyl,  $X_{10}$  is preferably substituted or unsubstituted furyl, thienyl, pyridyl, phenyl, or lower alkyl (e.g., tert-butyl),  $R_{2a}$  is preferably substituted or unsubstituted furyl, thienyl, pyridyl, phenyl, or lower alkyl, and  $R_7$ ,  $R_9$  and  $R_{10}$  each have the beta stereochemical configuration.

20 Any substituents of each  $X_3$ ,  $X_5$ ,  $R_2$ ,  $R_9$ ,  $R_{10}$  may be hydrocarbyl or any of the heteroatom containing substituents selected from the group consisting of heterocyclo, alkoxy, alkenoxy, alkynoxy, aryloxy, hydroxy, protected hydroxy, keto, acyloxy, nitro, amino, amido, thiol, ketal, acetal, ester and ether moieties, but not phosphorous containing moieties.



(15)



Express Mail No. EL 998648560 US

Series	X <sub>5</sub>	X <sub>3</sub>	R <sub>10</sub>	R <sub>2</sub>	R <sub>9</sub>	R <sub>14</sub>
A1	-COOX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	H
A2	-COX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	H
A3	-CONHX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	H
A4	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	H
A5	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	H
A6	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	H
A7	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	H
A8	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	H
A9	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	H
A10	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	H
A11	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	H
A12	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	H
B1	-COOX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	H
B2	-COX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	H
B3	-CONHX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	H
B4	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	H

Express Mail No. EL 998648560 US

B5	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	H
B6	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	H
B7	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	H
B8	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	H
B9	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	H
B10	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	H
B11	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	H
B12	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	H
C1	-COOX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	R <sub>9a</sub> COO-	H
C2	-COX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	R <sub>9a</sub> COO-	H
C3	-CONHX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	R <sub>9a</sub> COO-	H
C4	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	R <sub>9a</sub> COO-	H
C5	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	R <sub>9a</sub> COO-	H
C6	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	R <sub>9a</sub> COO-	H
C7	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	R <sub>9a</sub> COO-	H

Express Mail No. EL 998648560 US

C8	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	R <sub>9a</sub> COO-	H
C9	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	R <sub>9a</sub> COO-	H
C10	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	R <sub>9a</sub> COO-	H
C11	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	R <sub>9a</sub> COO-	H
C12	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	R <sub>9a</sub> COO-	H
D1	-COOX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	H
D2	-COX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	H
D3	-CONHX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	H
D4	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	H
D5	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	H
D6	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	H
D7	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	H
D8	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	H
D9	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	H
D10	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	H

Express Mail No. EL 998648560 US

D11	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	H
D12	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	H
E1	-COOX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	OH
E2	-COX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	OH
E3	-CONHX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	OH
E4	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	OH
E5	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	OH
E6	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	OH
E7	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	OH
E8	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	OH
E9	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	OH
E10	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	OH
E11	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	OH
E12	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	O	OH
F1	-COOX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	H
F2	-COX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	H
F3	-CONHX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	H

Express Mail No. EL 998648560 US

F4	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	H
F5	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	H
F6	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	H
F7	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	H
F8	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	H
F9	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	H
F10	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	H
F11	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	H
F12	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	H
G1	-COOX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	H
G2	-COX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	H
G3	-CONHX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	H
G4	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	H
G5	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	H
G6	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	H

Express Mail No. EL 998648560 US

5	G7	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	H
	G8	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	H
	G9	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	H
	G10	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	H
	G11	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	H
	G12	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	H
10	H1	-COOX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	OH
	H2	-COX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	OH
	H3	-CONHX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	OH
	H4	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	OH
	H5	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	OH
	H6	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	OH
15	H7	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	OH
	H8	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	OH
	H9	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	OH

Express Mail No. EL 998648560 US

H10	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	OH
H11	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	OH
H12	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	C <sub>6</sub> H <sub>5</sub> COO-	OH	OH
I1	-COOX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	OH
I2	-COX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	OH
I3	-CONHX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	OH
I4	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	OH
I5	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	OH
I6	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	OH
I7	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	OH
I8	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	OH
I9	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	OH
I10	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	OH
I11	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	OH
I12	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	O	OH
J1	-COOX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	OH

Express Mail No. EL 998648560 US

J2	-COX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	OH
J3	-CONHX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	OH
J4	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	OH
J5	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	OH
J6	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	OH
J7	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	OH
J8	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	OH
J9	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	OH
J10	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	OH
J11	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	OH
J12	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	OH	OH
K1	-COOX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	OH
K2	-COX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	OH
K3	-CONHX <sub>10</sub>	heterocyclo	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	OH
K4	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	OH
K5	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	OH



Express Mail No. EL 998648560 US

K6	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	OH
K7	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	OH
K8	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	OH
K9	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkenyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	OH
K10	-COOX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	OH
K11	-COX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	OH
K12	-CONHX <sub>10</sub>	optionally substituted C <sub>2</sub> to C <sub>8</sub> alkynyl	R <sub>10a</sub> COO-	R <sub>2a</sub> COO-	R <sub>9a</sub> COO-	OH

Example 5*In Vitro* cytotoxicity measured by the cell colony formation assay

Four hundred cells (HCT116) were plated in 60 mm Petri dishes containing 2.7 mL of medium (modified McCoy's 5a medium containing 10% fetal bovine serum and 100 units/mL penicillin and 100 g/mL streptomycin). The cells were incubated in a CO<sub>2</sub> incubator at 37 °C for 5 h for attachment to the bottom of Petri dishes. The compounds identified in Example 2 were made up fresh in medium at ten times the final concentration, and then 0.3 mL of this stock solution was added to the 2.7 mL of medium in the dish. The cells were then incubated with drugs for 72 h at 37 °C. At the end of incubation the drug-containing media were decanted, the dishes were rinsed with 4 mL of Hank's Balance Salt Solution (HBSS), 5 mL of fresh medium was added, and the dishes were returned to the incubator for colony formation. The cell colonies were counted using a colony counter after incubation for 7 days. Cell survival was calculated and the values of ID<sub>50</sub> (the drug concentration producing 50% inhibition of colony formation) were

Express Mail No. EL 998648560 US

determined for each tested compound.

Compound	IN VITRO ID 50 (nm) HCT116
taxol	2.1
docetaxel	0.6
0499	<1
0503	<1
0517	<10
0521	<1
0536	<1
0549	<10
0550	<10
0562	<1
0578	<1
0583	<10
0596	<10
0602	<1
0611	<10
0625	<1
0634	<10
0647	12.0
0659	<1
0663	<1
0670	<1
0687	<1
0691	<1
0706	<1
0719	<10
0720	<10

Express Mail No. EL 998648560 US

5

10

15

20

25

30

0732	<10
0748	<10
0838	<1
0843	<1
0854	<1
0860	<1
0879	<1
0882	<1
0890	<1
0908	<1
0919	<1
0923	<1
0937	<10
0947	<1
0951	<1
0966	<10
0978	<1
0983	<1
0999	<1
1003	<1
1011	<1
1020	<1
1031	<10
1044	<1
1060	<1
1879	<10
1883	<10
1892	<1
1900	<1
1911	<10

Express Mail No. EL 998648560 US

5

10

15

20

25

30

1923	<1
1939	<1
1948	<10
1954	<1
1964	<10
1970	<10
1988	<10
2101	<1
2111	<1
2124	<10
2132	<1
2142	<1
2159	<1
2164	<1
2173	<1
2181	<10
2199	<10
2202	<1
2212	<10
2226	<1
2238	<1
2242	<10
2255	<1
2269	<1
2273	<1
2287	<1
2291	<1
2306	<10
2319	<10
2320	<1

Express Mail No. EL 998648560 US

5

10

15

20

25

30

2332	<1
2348	<1
2353	<10
2366	<1
2379	<1
2380	<1
2392	<1
2408	<1
2413	<10
2424	<10
2439	<10
2442	<1
2455	<10
2464	<1
2472	<1
2488	<1
2499	<1
2503	<1
2511	<1
2520	<10
2781	<1
2794	<1
2802	<1
2813	<1
2826	<1
2838	<1
2844	<10
2855	<1
2869	<10
3053	<1

Express Mail No. EL 998648560 US

5

10

15

20

25

30

3071	<1
3096	<1
3102	<1
3110	<1
3129	<10
3132	<1
3148	<1
3163	<1
3204	<1
3219	<1
3222	<1
3258	<1
3265	<10
3297	<1
3314	<1
3352	<1
3361	<1
3370	<1
3408	<1
3417	<1
3425	<1
3453	<1
3482	<1
3494	<1
3513	<1
3522	<1
3535	<1
3543	<10
3588	<10
3595	<1

Express Mail No. EL 998648560 US

5

10

15

20

25

30

3603	<10
3644	<1
3656	<1
3663	<1
3677	<1
3686	<1
3693	<1
3800	<1
3818	<1
3853	<1
3866	<1
3909	<1
3938	<10
3945	<1
3957	<10
3971	<1
3982	<1
3994	<1
4051	<1
4062	<1
4112	<10
4121	<10
4190	<10
4207	<10
4329	<1
4335	<1
4344	<1
4665	<10
4704	<10
4711	<10

**Express Mail No. EL 998648560 US**

4720	<10
4799	<1
4808	<10
4834	<10
4888	<1
4919	<1
4944	<1
5011	<10
5040	<1
5065	<10
5144	<10
5232	<10
5495	<1
6522	<1

5

10



Express Mail No. EL 998648560 US

Example 6

## Preparation of Solutions for Oral Administration

5       Solution 1:   Antitumor compound 0499 was dissolved in ethanol to form a solution containing 106 mg of the compound per ml of solution. An equal volume of Cremophor® EL solution was added to the solution while stirring to form a solution containing 53 mg of compound 0499 per ml of solution. This solution was diluted using 9 parts by weight of saline to form a pharmaceutically acceptable solution for administration to a patient.

10       Solution 2:   Antitumor compound 0550 was dissolved in ethanol to form a solution containing 140 mg of the compound per ml of solution. An equal volume of Cremophor® EL solution was added to the solution while stirring to form a solution containing 70 mg of compound 0550 per ml of solution. This solution was diluted using 9 parts by weight of saline to form a pharmaceutically acceptable solution for administration to a patient.

15       Solution 3:   Antitumor compound 0611 was dissolved in ethanol to form a solution containing 150 mg of the compound per ml of solution. An equal volume of Cremophor® EL solution was added to the solution while stirring to form a solution containing 75 mg of compound 0611 per ml of solution. This solution was diluted using 9 parts by weight of saline to form a pharmaceutically acceptable solution for administration to a patient.

20       Solution 4:   Antitumor compound 0748 was dissolved in ethanol to form a solution containing 266 mg of the compound per ml of solution. An equal volume of Cremophor® EL solution was added to the solution while stirring to form a solution containing 133 mg of compound 0748 per ml of solution. This solution was diluted using 9 parts by weight of saline to form a pharmaceutically acceptable solution for administration to a patient.